MONTANA DEPARTMENT OF TRANSPORTATION WETLAND MITIGATION MONITORING REPORT: YEAR 2009

Big Hole Grazing Association Beaverhead County, Montana



Prepared for:



Prepared by:



POST, BUCKLEY, SCHUH, AND JERNIGAN 820 North Montana Avenue, Suite A Helena, MT 59601

December 2009

PBS&J Project No: 0B4308802.03.02

MONTANA DEPARTMENT OF TRANSPORTATION

WETLAND MITIGATION MONITORING REPORT

YEAR 2009

Big Hole Grazing Association Beaverhead County, Montana

MDT Project Number STPX 1(45) Control Number 4668

Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION

2701 Prospect Ave Helena, MT 59620-1001

Prepared by:

POST, BUCKLEY, SCHUH, AND JERNIGAN 820 North Montana Avenue, Suite A Helena, MT 59601

December 2009

PBS&J Project No: 0B4308802.03.02

"MDT attempts to provide accommodations for any known disability that may interfere with a person participating in any service, program, or activity of the Department of Transportation. Alternative accessible formats of this information will be provided upon request. For further information, call 406-444-7228, TTY at 800-335-7592, or Montana Relay at 711."

TABLE OF CONTENTS

1.0	INTRODUCTION1
2.0	METHODS3
	2.1 Monitoring Dates and Activities
	2.2 Hydrology
	2.3 Vegetation
	2.4 Soils
	2.5 Wetland Delineation4
	2.6 Mammals, Reptiles, and Amphibians4
	2.7 Birds
	2.8 Macroinvertebrates5
	2.9 Functional Assessment5
	2.10 Photographs5
	2.11 GPS Data5
	2.12 Maintenance Needs
3.0	RESULTS5
	3.1 Hydrology5
	3.2 Vegetation
	3.3 Soils9
	3.4 Wetland Delineation
	3.5 Wildlife
	3.6 Macroinvertebrates
	3.7 Functional Assessment
	3.8 Photographs
	3.9 Maintenance Needs/Recommendations
	3.10 Current Credit Summary
4.0	REFERENCES14



TABLES

Table 1 Vegetation species observed from 2008 to 2009 at the Big Hole Grazing Association Wetland Mitigation Site.
 Table 2 Data summary from 2008 to 2009 for Transect 1.
 Table 3 Acreages for each wetland community from 2008 to 2009 at the Big Hole Grazing Association Wetland Mitigation Site.
 Table 4 Fish and wildlife species observed within the Big Hole Grazing Association Wetland Mitigation Site in 2008 and 2009.
 Table 5 Summary of 2009 wetland function/value ratings and functional points at the Big Hole Grazing Association Wetland Mitigation Site.

FIGURES

Figure 1 Project Site Location Map
Figure 2 2009 Monitoring Activity Locations
Figure 3 2009 Mapped Site Features

CHARTS

Chart 1 Transect map showing vegetation types of Transect 1 from start (0 feet) to end (1247 feet) in 2008 and 2009.

Chart 2 Length of vegetation communities within Transect 1 during 2008 and 2009.

APPENDICES

Appendix A Figures 2 & 3

Appendix B 2009 Wetland Mitigation Site Monitoring Form

2009 Bird Survey Form

2009 COE Wetland Delineation Forms 2009 MDT Functional Assessment Forms

Appendix C 2009 Representative Photographs

Appendix D Project Plan Sheet

Appendix E Bird Survey Protocol

GPS Protocol



1.0 INTRODUCTION

The Big Hole Grazing Association (BHGA) wetland mitigation project was constructed in the fall of 2007 by the Montana Department of Transportation (MDT). The purpose of the project is to restore approximately 45 acres of wetland habitat within a 96-acre easement area owned by the BHGA. The project would provide a wetland mitigation reserve in Watershed #6 - Upper Missouri River Basin. MDT has not yet identified any construction projects in this watershed that would be applied against the Big Hole Grazing Association reserve.


This report documents the second year of monitoring at the Big Hole Grazing Association Wetland Mitigation site. This project is located approximately seven miles southwest of the town of Wisdom and approximately four miles west of Secondary Route 278 (**Figure 1**). Specifically, the project is located in the NW ¼ of Section 2, Township 4 South and Range 16 West in Beaverhead County.

Prior to project initiation, the BHGA utilized the project area for grazing and haying operations. Drainage ditches had been utilized for decades to drain the project area, which has a naturally high groundwater table and receives spring flows from the hillside to the north of the site. In addition to the springs and groundwater, Rock Creek, a perennial tributary to the Big Hole River, flows through the southern portion of the easement area.

In an attempt to restore natural hydrology to the site and thus restore wetlands within the easement area, the primary drainage ditch, which flowed northwest to southeast through the easement area, was completely filled and reclaimed. A secondary ditch which runs north to south across the western portion of the site was breached in three locations in an attempt to prevent drainage of the site and re-hydrate former wetlands by raising groundwater levels across the site. A project plan sheet is provided in **Appendix D**.

Prior to project implementation, MDT documented approximately 31 acres of degraded and relic emergent and scrub/shrub wetland across the entire 96-acre easement area, noting that some wetland areas were likely much larger prior to construction of drainage ditches across the site in the 1960's. The intent of the project was to restore natural hydrology to the site in an attempt to restore wetlands within the easement area. According to project files, the goal is to generate 45.8 acres of Corp of Engineers approved credit through the restoration of 42.3 acres of wetland credited at 1:1 and preservation of 14.0 acres credited at a 4:1 ratio (3.5 acres of credit).





2.0 METHODS

2.1 Monitoring Dates and Activities

Monitoring in 2009 occurred on August 12th. All information contained on the Wetland Mitigation Site Monitoring Form was collected during this site visit (**Appendix B**). Activities conducted and information collected included: wetland delineation; vegetation community mapping; vegetation transect monitoring; soils data collection; hydrology data collection; mapping of man-made bird nest boxes; bird and wildlife use documentation; and photo documentation.

2.2 Hydrology

Hydrologic indicators were evaluated during the site visit on August 12th. Wetland hydrology indicators were recorded using procedures outlined in the COE 1987 Wetland Delineation Manual (Environmental Laboratory 1987). Hydrology data were recorded on COE Routine Wetland Delineation Data Forms and on the Wetland Mitigation Site Monitoring Form (**Appendix B**).

There are eight groundwater monitoring wells at the site that, up until 2009, were routinely monitored by the U.S. Geological Service (USGS). The USGS discontinued monitoring of the wells in 2009 at the request of MDT and the monitoring wells were not measured by PBS&J during the site visit. Soil pits excavated for wetland delineation purposes were used to evaluate the presence of groundwater if occurring within 12 inches from the ground surface. Data were recorded on the COE Routine Wetland Delineation Data Form (**Appendix B**)

2.3 Vegetation

General dominant species-based vegetation community types were delineated in the field during the mid-summer field visit. Standardized community mapping was not employed as many of these systems are geared towards climax vegetation. Estimated percent cover of the dominant species in each community type was recorded on the Wetland Mitigation Site Monitoring Form (**Appendix B**). Plants observed were identified using Flora of the Pacific Northwest (Hitchcock and Conquist 1975) and Plants of Montana (Dorn 1984). Nomenclature follows that of Dorn (1984).

Annual changes in vegetation, especially the establishment and increase of hydrophytic plants, will be evaluated through the use of belt transects. A single 10-foot wide vegetation belt transect was established at the site and monitored for the second time in 2009. The transect start and end points were marked in the field and recorded with a global positioning system (GPS) unit in 2008. Percent cover was estimated for each successive vegetative species encountered within the "belt" using the following values: +(<1%); 1 (1-5%); 2 (6-10%); 3 (11-20%); 4 (21-50%); and 5 (>50%). Photographs were taken at the start and end of the transect during the mid-season visit.



3

Woody species were planted in clumps across the site in May 2008. The location of each clump was recorded in the field with GPS in 2008 and examined for plant survival for the second time in 2009.

2.4 Soils

Soil information was obtained from the Web Soil Survey (NRCS 2008). Soils were evaluated during the mid-season visit according to procedures outlined in the COE 1987 Wetland Delineation Manual. In the field, surface soils were evaluated for signs of wetland formation during the mid-season visit. If wetland indicators for hydrology or plants were found then a soil pit was excavated to look for evidence of hydric soil formation. Soil data were then recorded on the COE Routine Wetland Delineation Form (**Appendix B**).

2.5 Wetland Delineation

Wetland delineation was conducted during the mid-season visit in accordance with the 1987 COE Wetland Delineation Manual. In July 2008, consultation with the COE (Steinle pers. comm.) confirmed that, where the 1987 manual was used to establish baseline wetland conditions at MDT wetland mitigation sites, it should continue to be applied at such sites for the duration of the monitoring period. Consequently, application of the new *Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Western Mountains*, *Valleys, and Coast Region* (COE 2008) was not required or undertaken at this site in 2008 or 2009.

The monitoring area was investigated for the presence of wetland hydrology, hydrophytic vegetation, and hydric soils. The indicator status of vegetation was derived from the National List of Plant Species that Occur in Wetlands: Northwest Region 9 (Reed 1988). The information was recorded on a COE Routine Wetland Delineation Data Form (**Appendix B**).

2.6 Mammals, Reptiles, and Amphibians

Mammal, reptile, and amphibian species observations and other positive indicators of use, such as vocalizations, were recorded on the wetland monitoring form during the site visit. Indirect use indicators, including tracks, scat, burrow, eggshells, skins, and bones, were also recorded. These signs were recorded as the observer traversed the site while conducting other required activities. Direct sampling methods, such as snap traps, live traps, and pitfall traps, were not used. A comprehensive wildlife species list for the entire site was compiled (**Appendix B**).

2.7 Birds

Bird observations were recorded during the site visit. No formal census plots, spot mapping, point counts, or strip transects were conducted. Bird observations were recorded incidental to other monitoring activity observations, using the bird survey protocol as a general guideline (**Appendix E**). Observations were categorized by species, activity code, and general habitat association on the Bird Survey Field Data Sheet (**Appendix B**). Shortly after monitoring was completed in 2008, MDT installed blue bird and wood duck nesting boxes across the site. Each



of these boxes was examined for use in 2009. A comprehensive bird list was compiled using these observations.

2.8 Macroinvertebrates

Due to the lack of standing water, no aquatic macroinvertebrate sample was collected at this site.

2.9 Functional Assessment

In 2001, a functional assessment for each delineated wetland was conducted using the 1999 MDT Montana Wetland Assessment Method (MWAM) (Berglund 1999). In 2009 the 2008 MDT Montana Wetland Assessment Method (Berglund and McEldowney 2008) was applied. Field data necessary for this assessment were collected during the mid-season site visit. For each wetland or group of wetlands a Functional Assessment Form was completed (**Appendix B**).

2.10 Photographs

Photographs were taken in 2009 to show the current land use surrounding the site, the upland buffer, the monitored area, and the vegetation transect. Seven photograph points were established and their location recorded with a resource grade GPS unit in 2008. A description and compass direction for each photograph was recorded onto the Wetland Mitigation Site Monitoring Form (**Appendix B**).

2.11 GPS Data

During the 2009 monitoring season, survey points were collected with a resource grade GPS unit at each of the bird boxes installed by MDT in 2008. Rather than using the GPS unit to survey all wetland boundaries within project limits, minor changes in wetland boundaries were hand drawn onto aerial photography in 2009 (**Appendix E**).

2.12 Maintenance Needs

The Big Hole Grazing Association mitigation site is a groundwater driven project that does not include any manmade diversions, water level control structures, or other structures that might need periodic maintenance.

3.0 RESULTS

3.1 Hydrology

Eight groundwater monitoring wells were installed across the site in 2001 and were monitored annually by the USGS through 2008. The primary goal of the project was to raise groundwater levels across the easement area by plugging the two main drainage ditches across the site. As indicated in the 2008 monitoring report, groundwater levels following construction in 2007 were noticeably higher during monitoring in 2008.



For the second time in 2009, water levels were documented (visual observation) above the ground surface at wells 6, 7, and 8. Although monitoring of the wells was discontinued in 2009, saturation levels and evidence of shallow standing surface water in several wet meadow locations indicate that groundwater levels were comparable in 2009 to those documented in 2008.

From January to September of 2009, the Wisdom weather station (#249067) reported 10.25 inches (in) of precipitation compared to the long term average of 9.53 inches over the same 9-month time period (Western Regional Climate Center (WRCC) 2009). It was assumed that precipitation levels measured at the Wisdom Weather Station would serve as an indicator of precipitation received at the mitigation site.

3.2 Vegetation

The relatively flat 96-acre easement area has a variety of wetland and upland vegetation communities. Plant species observed during the first two years of monitoring have been compiled in a list (**Table 1**). Mapped vegetation community types were based on topography, hydrology, and plant composition. A total of one upland community and five wetland communities were identified in 2009 (**Figure 3** in **Appendix A**). Transitional wetland identified in 2008 was reclassified as a combination of wetland communities 3 and 5 in 2009. For purposes of describing vegetation types, the site can be divided into four basic areas: the northwest corner, the northeast corner, the south side, and the central portion. The northwest corner contains a 7-acre sedge-dominated (Vegetation Type 3) fen that is in reasonably good health and was likely never adversely impacted by previous ditching on the site (**Monitoring Forms** in **Appendix B**).

The east/northeast side of the easement area was once likely dominated by willow communities but was converted to upland and wet meadow through years of being drained and grazed. This area has numerous hummocks and with the restoration of hydrology to the site is already showing signs of converting back to its previous state. Willow shoots are common on the numerous hummocks, while sedge and other OBL and FACW species are replacing more facultative grasses and forbs (Vegetation Type 6) (Monitoring Forms in Appendix B).

Rock Creek and its associated willow-dominated (Vegetation Type 4) floodplain run west to east across the southern portion of the easement area (**Monitoring Forms** in **Appendix B**). Willow communities are in relatively good health and were never adversely affected by drainage of the easement area because these wetland areas are tied hydraulically to Rock Creek.

The large central portion of the easement area is a mix of Vegetation Type 1 (mixed grass upland) and Type 5 (wet meadow) (**Monitoring Forms** in **Appendix B**). This area has been most affected by past management activities on the property, as wetland was converted to upland for agricultural purposes.



Table 1: Vegetation species observed from 2008 to 2009 at the Big Hole Grazing Association

Wetland Mitigation Site.

Scientific Name	Region 9 (Northwest) Wetland Indicator	Scientific Name	Region 9 (Northwest) Wetland Indicator
Achillea millefolium	FACU	Hordeum jubatum	FAC+
Agropyron trachycaulum	FAC	Juncus balticus	OBL
Agrostis alba	FACW	Juncus longistylis	FACW
Alnus incana	FACW	Kochia scoparia	FAC
Alopecurus pratensis	FACW	Lemna minor	OBL
Aster hesperius	OBL	Mentha arvense	FAC
Beckmannia syzigachne	OBL	Mimulus guttatus	OBL
Betula pumila	OBL	Myosotis scorpioides	FACW
Bromus inermis		Phleum pratense	FACU
Calamagrostis canadensis	FACW+	Poa juncifolia	FACU+
Carex aquatilis	OBL	Poa pratensis	FACU+
Carex nebrascensis	OBL	Polygonum amphibium	OBL
Carex praegracilis	FACW	Potamogeton sp.	OBL
Carex utriculata	OBL	Potentilla glandulosa	FAC-
Centaurea maculosa		Potentilla gracilis	FAC
Cirsium arvense	FACU+	Rosa woodsii	FACU
Cornus stolonifera	FACW	Rumex crispus	FACW
Deschampsia cespitosa	FACW	Salix bebbiana	FACW
Eleocharis palustris	OBL	Salix exigua	OBL
Eleocharis pauciflora	OBL	Salix lemmonii	FACW+
Epilobium ciliatum	FACW-	Scirpus acutus	OBL
Equisetum arvense	FAC	Senecio sphaerocephalus	FACW
Festuca octoflora		Sisyrinchium angustifolium	FACW-
Geum Aleppicum	FACW-	Spiranthes romanzoffiana	OBL
Glyceria elata	FACW+	Taraxacum officinale	
Glycyrrhiza lepidota	FAC+	Thlaspi arvense	
Gnaphalium palustre	FAC+	Trifolium pratense	FACU
Hippuris vulgaris	OBL	Triglochin palustre	OBL
Hordeum brachyantherum	FACW-	Typha latifolia	OBL

Vegetation Type 3/5 occurs along the large drainage ditch that used to drain the project area but was filled as part of the restoration effort. Wetland vegetation is beginning to establish in this area as a result of seeding and planting of woody vegetation (**Monitoring Forms** in **Appendix B**). This area was identified as transitional wetland in 2008, but due to extensive wetland vegetation establishment in 2009, was reclassified.


Plant composition along a single vegetation transect was quantified during the monitoring effort in 2009. Transect results are detailed in the Monitoring Form (**Appendix B**) and are summarized in **Table 2** and **Charts 1** and **2**. The transect runs south to north through the central portion of the mitigation area; it starts at monitoring well #3 and ends at monitoring well #6 (**Figure 2** in **Appendix A**). This transect includes areas of mixed grass upland, wet meadow, and developing emergent marsh where the drainage ditch was filled.



Table 2: Data summary from 2008 to 2009 for Transect 1.

Monitoring Year	2008	2009
Transect Length (feet)	1247	1247
# Vegetation Community Transitions along Transect	7	7
# Vegetation Communities along Transect	3	3
# Hydrophytic Vegetation Communities along Transect	2	2
Total Vegetative Species	22	22
Total Hydrophytic Species	13	14
Total Upland Species	9	8
Estimated % Total Vegetative Cover	75	80
% Transect Length Comprised of Hydrophytic Vegetation Communities	45	55
% Transect Length Comprised of Upland Vegetation Communities	55	45
% Transect Length Comprised of Unvegetated Open Water	0	0
% Transect Length Comprised of Bare Substrate	0	0

Chart 1: Transect map showing vegetation types of Transect 1 from start (0 feet) to end (1247 feet) in 2008 and 2009.








Chart 2: Length of vegetation communities within Transect 1 during 2008 and 2009.

In May 2008, woody vegetation was planted in 45 clumps along the filled-in drainage ditch and along the secondary drainage ditch that was plugged in three locations (**Figure 2** in **Appendix A**). Three plant species were planted: birch (*Betula pumila*), alder (*Alnus incana*), and dogwood (*Cornus stolonifera*). There were a total of 961 plantings counted during the 2008 monitoring. Approximately 79% (756 stems) survived through the first growing season. In 2009, the second year of monitoring, survival dropped considerably to 35% (339 stems). Of the three species planted, alder is faring the best, at 55% survival, while red-osier dogwood survival is about 30% and birch less than 2%. Mortality of birch and dogwood plantings was likely due to wetter than expected conditions and competition from forbs and grasses. It may also be possible that these species are not suited for the environment within which they were planted.

Cirsium arvense was the only State noxious weed found in 2009; it occurred in two primary locations near each other (**Figure 3** in **Appendix A**). The infestation is small, but appeared to have expanded slightly from 2008.

3.3 Soils

According to soil survey, the easement area has two primary soil types: *Mooseflat Loam 0-4% slope* along the Rock Creek corridor and Foxgulch-Copperbasin-Wisdom complex 0-2% slope for the remaining study area (NRCS 2008). The drainage class for both soil types is conducive to wetland development.



Soils were investigated across much of the site in 2009. Typical soil profiles throughout the site consisted of sandy loam and clay loam with matrix colors ranging from 10YR 2/1 with no mottles to 10YR 4/2 with distinct 10YR 6/8 mottles (**COE Forms** in **Appendix B**). Within wetland areas, soils were generally saturated within the upper 12 inches of the profile and to near the surface in many cases.

3.4 Wetland Delineation

According to project documents, the site was delineated by MDT in June of 2001. At that time, approximately 31 acres of drained and degraded wetland habitat was delineated within project boundaries. Wetland delineation during the August 2008 monitoring documented 49.81 acres of wetland, which increased to 56.76 acres in 2009 for a net gain of 25.76 acres since project inception (**Figure 3** in **Appendix A**). The largest difference between the 2001 delineation and 2009 delineation is in the east/northeast corner, where the 2009 delineation revealed 13.45 acres of developing EM/SS wetland in an area previously determined by MDT to be upland. It is probable that restoration of groundwater to this part of the project in the fall of 2007 was already having a positive effect on plant composition and saturated soil conditions in August 2009.

Delineated wetland boundaries, uplands, and transitional areas were mapped (**Figure 3** in **Appendix A**). As mentioned, hydrophytic vegetation was quick to establish in the northeast corner of the site and also began to appear within the central portion of the site in previously identified upland areas. Herbaceous plant establishment in the filled drain ditch expanded significantly in 2009, as disturbed areas transition to wetland (Wetland Type 3/5 on **Figure 3** in **Appendix A**).

Acreages were calculated for delineated wetlands within the monitoring limits (**Table 3**). In addition to the 56.76 acres of wetland within monitoring limits, there is an additional approximately 38.1 acres of upland habitat.

Table 3: Acreages for each wetland community from 2008 to 2009 at the Big Hole Grazing Association Wetland Mitigation Site.

WETLAND COMMUNITY	ACREAGE	
WEILAND COMMUNITY	2008	2009
Type 3 – <i>Carex</i> Wetland	7.19	7.19
Type 4 – Salix Wetland	13.28	13.28
Type 5 – Wet Meadow Wetland	15.89	21.20
Type 3/5 – Developing Emergent / Wet Meadow Wetland	0.00	1.64
Type 6 – Emergent / Scrub-shrub Wetland	13.45	13.45
Total Wetland Habitat	49.81	56.76

3.5 Wildlife

Direct observations of all wildlife species and signs indicating their presence were compiled in **Table 4** and in **Appendix B**. Three moose were seen on the site during the field survey and according to the land owner, a cow and calf are observed regularly. Relatively few bird species were noted – however, conditions were less than optimal (temperatures in the 80's) during the



10

August monitoring. Of note was the observation of a northern harrier family group. The land owner documented a harrier nest on the property earlier in the summer. Animals not observed but likely to occur onsite include various small mammals, reptiles, amphibians, and various song birds.

Table 4: Fish and wildlife species observed within the Big Hole Grazing Association Wetland Mitigation Site in 2008 and 2009.

Miligation Site in 2008 and 2009.	
FISH, AMPHIBIAN, and REPTILE	
Trout (unidentified species in Rock Creek)	
BIRD	
Bank Swallow (Riparia riparia) Cliff Swallow (Petrochelidon pyrrhonota) Common Raven (Corvus corax) Common Snipe (Gallinago gallinago) Great Horned Owl (Bubo virginianus) Great Blue Heron (Ardea herodias) Killdeer (Charadrius vociferous) Black-billed Magpie (Pica pica) Mallard (Anas platyrhynchos)	Northern Harrier (Circus cyaneus) Red-winged Blackbird (Agelaius phoeniceus) Rock Dove (Columba livia) Sandhill Crane (Grus Canadensis) Spotted Sandpiper (Actitis macularia) Tree Swallow (Tachycineta bicolor) Wilson's Phalarope (Phalaropus tricolor) Song Sparrow (Melospiza melodia)
MAMMAL	
Deer (Odocoileus sp.) Moose (Alces alces)	American Badger (Taxidea taxus)

Bolded species were observed during the 2009 monitoring year while other species were observed in previous monitoring years.

3.6 Macroinvertebrates

Aquatic macroinvertebrates were not sampled at this site in 2009 because the site did not contain large areas of surface water.

3.7 Functional Assessment

MDT project files indicate that prior to project implementation, wetlands occurring along the Rock Creek corridor and in the northwest corner (fen area) rated as Category II wetlands while all other wetland on the site rated as Category III using the MDT 1999 MDT Montana Wetland Assessment Method. The 2009 conditions were assessed using the 2008 Montana Wetlands Assessment Method (Functional Assessment Forms in Appendix B).

In 2009, two functional assessments were completed for wetlands at the Big Hole Grazing Association mitigation site. Wetlands associated with the Rock Creek drainage (Assessment 1) were assessed on one form while all other wetlands were rated together (Assessment 2). In summary, all wetlands within project boundaries rated as Category II, with high ratings for short and long term surface water storage, sediment/nutrient/toxicant removal, and groundwater discharge/recharge (**Table 5**). Wetlands along Rock Creek also received high ratings for fish and wildlife habitat, flood attenuation, short and long term surface water storage, sediment/



shoreline protection, and production export / food chain support. The larger wetland area received a high rating for uniqueness because of the fen qualities exhibited in the northwest portion of the easement area.

Table 5: Summary of 2009 wetland function/value ratings and functional points at the Big

Hole Grazing Association Wetland Mitigation Site.

Function and Value Parameters from the	2009	2009
MDT Montana Wetland Assessment	Assessment 1	Assessment 2
Method	(Rock Creek Wetlands)	(Remaining Wetlands)
Listed/Proposed T&E Species Habitat	Low (0.0)	Low (0.0)
MTNHP Species Habitat	Low (0.2)	Low (0.2)
General Wildlife Habitat	High (0.9)	Mod (0.7)
General Fish/Aquatic Habitat	High (0.8)	NA
Flood Attenuation	High (0.8)	NA
Short and Long Term Surface Water Storage	High (0.8)	High (1.0)
Sediment/Nutrient/Toxicant Removal	High (0.9)	High (1.0)
Sediment/Shoreline Stabilization	High (1.0)	NA
Production Export/Food Chain Support	High (1.0)	Mod (0.6)
Groundwater Discharge/Recharge	High (1.0)	High (1.0)
Uniqueness	Mod (0.4)	High (0.9)
Recreation/Education Potential	Low(0.05)	Low(0.05)
Actual Points / Possible Points	7.85 / 11	5.45 / 8
% of Possible Score Achieved	71%	68%
Overall Category	II	II
Total Acreage of Assessed Wetlands and		
Other Aquatic Habitats within Site	10	39.81
Boundaries (ac)		
Functional Units (acreage x actual points)	78.50	217

3.8 Photographs

Representative photos were taken from photo-points (**Photos 1-21**), transect ends (**Photos 22-23**), and of the general project area (**Photo 24**) (**Appendix C**). The 2009 aerial photograph taken on July 1st was used as a base for **Figures 2** and **3** (**Appendix A**).

3.9 Maintenance Needs / Recommendations

As a majority of the mitigation site relies entirely on groundwater to support wetlands, there are no man-made water level control features to monitor. The project area has a wooden fence around the perimeter that was in good condition in 2009. All man-made bird nesting structures installed in 2008 by MDT were in good condition.

From a vegetative standpoint, disturbed areas where the ditch was filled were reseeded following construction and appeared to be progressing satisfactorily. Woody planting survival declined significantly in 2009 and may need to be supplemented in future years if desired by MDT. No additional seeding is recommended at this time. Two small infestation of Canada thistle located close together were identified (**Figure 3** in **Appendix A**). Spraying or pulling may prevent spread to other parts of the site.



3.10 Current Credit Summary

Correspondence in the MDT project file indicates that the overall goal was to achieve 45.8 acres of Corps-approved mitigation credit within the 96 acre easement area. Credit was to be obtained for 42.3 acres of wetland restoration at a ratio of 1:1, and 3.5 acres of credit was to be obtained for preservation of 14.0 acres at a ratio of 4:1. The 14.0 acres of preservation applies to the Rock Creek corridor and the fen area in the northwest corner of the site – neither of which was impacted by the filling of the ditches. These areas did however benefit from the removal of cattle grazing from the site.

Assuming that restoration credit is granted for all wetlands on the site other than the 14 acres dedicated as "preservation", the maximum 2009 credit breakdown is as follows:

14 acres preservation at 4:1 ratio:
42.76 acres restoration at 1:1 ratio:
42.76 acres credit
42.76 acres credit
42.76 acres credit
46.26 acres credit

Years 1 and 2 of monitoring have shown a positive trend toward hydrologic restoration within a large portion of the site as indicated by the fact that 56.76 acres of wetland were delineated on the site compared to 31 acres pre-project. Over time, it is anticipated that additional wetland will develop, especially towards the center of the project area, where upland grass communities may convert to wet meadow and emergent marsh. After year two of monitoring, the site has already exceeded the intended goal of 45.8 acres of credit. Additional acreage is possible and will be documented during future monitoring efforts.



4.0 REFERENCES


- Berglund, J. 1999. *MDT Montana Wetland Assessment Method*. May 25th. Prepared for Montana Department of Transportation and Morrison-Maierle, Inc. Prepared by Western EcoTech. Helena, Montana. 18 pp.
- Berglund, J. and R. McEldowney. 2008. *MDT Montana Wetland Assessment Method*. Prepared for Montana Department of Transportation, Helena, Montana. Post, Buckley, Schuh, & Jernigan, Helena, Montana. 42pp.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. U.S. Army Corps of Engineers. Washington, DC.
- Natural Resources Conservation Service (NRCS). 2008. Soil Information for the Big Hole Grazing Wetland Mitigation Site. Obtained on December 8th from the world wide web at: http://websoilsurvey.nrcs.usda.gov/app/.
- Reed, P.B. 1988. *National List of Plant Species that Occur in Wetlands: North West (Region 9)*. Biological Report 88(26.9), May 1988. U.S. Fish and Wildlife Service. Washington, D.C.
- Steinle, A. 2008. Montana Program Manager, U.S. Army Corps of Engineers, Helena, Montana. July 14th telephone conversation.
- U.S. Army Corps of Engineers (COE). 2008. *Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region*, ed. J. S. Wakeley, R. W. Lichvar, and C. V. Noble. ERDC/EL TR-08-13. U.S. Army Engineer Research and Development Center, Vicksburg, Missouri.
- Western Regional Climate Center (WRCC). 2009. Precipitation data for the Wisdom weather station, Montana (249067). Obtained on October 30, 2009 from http://www.wrcc.dri.edu/CLIMATEDATA.html




Appendix A

FIGURES 2 & 3

MDT Wetland Mitigation Monitoring Big Hole Grazing Association Beaverhead County, Montana





Appendix B

2009 WETLAND MITIGATION SITE MONITORING FORM 2009 BIRD SURVEY FORM 2009 COE WETLAND DELINEATION FORMS 2009 MDT FUNCTIONAL ASSESSMENT FORMS

MDT Wetland Mitigation Monitoring Big Hole Grazing Association Beaverhead County, Montana

PBS&J/MDT WETLAND MITIGATION SITE MONITORING FORM

Project Name: <u>Bighole Grazing Association</u> Project Number: <u>0B4308802</u> Assessment Date: <u>August 12, 2009</u> Person(s) conducting the assessment: <u>Traxler</u> Location: <u>10 miles southwest of Wisdom</u> MDT District: <u>Butte</u> Milepost: Legal Description: T <u>4S</u> R <u>16W Section 2</u> Weather Conditions: <u>partly cloudy, warm</u> Time of Day: <u>7:30am - 1:00pm</u> Initial Evaluation Date: <u>August 6, 2008</u> Monitoring Year: <u>2</u> # Visits in Year: <u>1</u> Size of evaluation area: <u>95 acres</u> Land use surrounding wetland: <u>Rangeland; hayland</u>					
HYDROLOGY					
Surface Water Source: Rock Creek, precipitation, springs, groundwater Inundation: Present Average Depth: Range of Depths: 0-36" Percent of assessment area under inundation: 67% Depth at emergent vegetation-open water boundary: 0.5 feet If assessment area is not inundated then are the soils saturated within 12 inches of surface: Yes Other evidence of hydrology on the site (ex. – drift lines, erosion, stained vegetation, etc.): Ditches and drainage patterns.					
Groundwater Monitoring Wells: Present Record depth of water below ground surface (in feet): not recorded in 2009					
Well Number Depth Well Number Depth Well Number Depth					
1 5					
$egin{array}{ c c c c c c c c c c c c c c c c c c c$					
3 7 4 8					
Additional Activities Checklist: Map emergent vegetation-open water boundary on aerial photograph. Observe extent of surface water during each site visit and look for evidence of past surface water elevations (drift lines, erosion, vegetation staining, etc.) Use GPS to survey groundwater monitoring well locations, if present. COMMENTS / PROBLEMS: USGS discontinued monitoring groundwater levels at the 8 well sites in 2009. Well readings were not taken by PBS&J during the August monitoring. As was the case in 2008, surface water was noted at well numbers 6 & 7. Based on visual observations of surface water and overall ground saturation across the site it is thought that groundwater levels were at least equal to or higher than					

the first year of monitoring (2008).

VEGETATION COMMUNITIES

Community Number: **1** Community Title (main spp): **Mixed Grass Upland**

Dominant Species	% Cover	Dominant Species	% Cover
Achellea millefolium	1 = 1-5%	Poa juncifolia	1 = 1-5%
Agropyron trachycaulum	2 = 6-10%	Poa pratensis	2 = 6-10%
Agrostis alba	2 = 6-10%	Taraxacum officinale	1 = 1-5%
Bromus inermis	1 = 1-5%		
Festuca octoflora	4 = 21-50%		
Phleum pratense	3 = 11-20%		

Comments / Problems: _____

Community Number: 2 Community Title (main spp): Transitional

TIP				
Dominant Species	% Cover	Dominant Species	% Cover	
Agrostis alba	2 = 6-10%	Juncus balticus	2 = 6-10%	
Beckmannia syzigachne	1 = 1-5%	Polygonum amphibium	1 = 1-5%	
Carex nebrascensis	1 = 1-5%	Potamogeton sp.	2 = 6-10%	
Eleocharis palustris	2 = 6-10%	Rumex crispus	1 = 1-5%	
Hordeum brachyantherum	2 = 6-10%	Mimulus guttatus	2 = 6-10%	
Hordeum jubatum	2 = 6-10%	Deschampsia cespitosa	2 = 6-10%	

Comments / Problems: Area where main drain ditch was filled in. Not present in 2009. This area developed significant wetland characteristics in 2009 and was classified as Community number 3/5.

Community Number: 3 Community Title (main spp): Carex

Dominant Species	% Cover	Dominant Species	% Cover
Carex aquatilis	4 = 21-50%		
Carex nebrascensis	4 = 21-50%		
Carex utriculata	4 = 21-50%		

Comments / Problems: Sedge dominated fen area in NW corner of site.

Community Number: 4 Community Title (main spp): Salix

community rumber. 4 Community rice (main spp). Dank				
Dominant Species	% Cover	Dominant Species	% Cover	
Salix exigua	3 = 11-20%	Agrostis alba	1 = 1-5%	
Salix lemmonii	3 = 11-20%	Juncus balticus	1 = 1-5%	
Salix bebbiana	3 = 11-20%	Calamagrostis canadensis	1 = 1-5%	
Alnus incana	2 = 6-10%			
Betula pumila	1 = 1-5%			
Cornus stolonifera	1 = 1-5%			

Comments / Problems: <u>Dominated by willow with scattered birch, dogwood, and alder in overstory and various wetland grasses in understory.</u>

VEGETATION COMMUNITIES (continued)

Community Number: <u>5</u> Community Title (main spp): <u>Wet Meadow (mixed herbaceous)</u>

Dominant Species	% Cover	Dominant Species	% Cover
Agrostis alba	2 = 6-10%	Deschampsia cespitosa	2 = 6-10%
Eleocharis palustris	1 = 1-5%	Mimulus guttatus	1 = 1-5%
Hordeum brachyantherum	2 = 6-10%	Senecio sphaerocephalus	1 = 1-5%
Hordeum jubatum	2 = 6-10%	Glyceria elata	2 = 6-10%
Juneus balticus	3 = 11-20%	Beckmannia syzigachne	2 = 6-10%
Alopecurus pratensis	3 = 11-20%		

Comments / Problems: Largest wetland community type within monitoring limits.

Community Number: 6 Community Title (main spp): EM/SS Mix

Dominant Species	% Cover	Dominant Species	% Cover
Agrostis alba	2 = 6-10%	Salix bebbiana	2 = 6-10%
Juneus balticus	3 = 11-20%	Salix lemmonii	2 = 6-10%
Carex nebrascensis	2 = 6-10%		
Carex utriculata	2 = 6-10%		
Carex praegracilis	1 = 1-5%		
Salix exigua	2 = 6-10%		

Comments / Problems: <u>Along eastern edge of site - shrub component is beginning to be restored after having grazing removed.</u>

Community Number: ___ Community Title (main spp):

Dominant Species	% Cover	Dominant Species	% Cover

Comments / Problems: _____

Community Number: ___ Community Title (main spp): ____

Dominant Species	% Cover	Dominant Species	% Cover

(Comments	/ Problems:	
ı	OHILLENIS	/ Problems	

COMPREHENSIVE VEGETATION LIST

Plant Species	Vegetation Community Number (s)	Plant Species	Vegetation Community Number (s)
Achillea millefolium	1,2	Juncus balticus	1,2,4,5,6
Agropyron trachycaulum	1	Juncus longistylus	1,2,5
Agrostis alba	1,2,4,5,6	Kochia scoparia	1
Alnus incana	4	Lemna minor	3,5,6
Glycyrrhiza lepidota	1,5	Mentha arvense	5
Alopecurus pratensis	1,2,5	Mimulus guttatus	5,6
Aster hesperius	3,5,6	Myosotis scorpioides	2,5,6
Beckmannia syzigachne	2,5	Phleum pratense	1
Betula pumila	4,6	Poa Juncifolia	1
Bromus inermis	1	Poa pratensis	1
Calamagrostis canadensis	3,4,5,6	Polygonum amphibium	2,6
Calamagrostis scopulorum	5,6	Potamogeton sp.	3,5,6
Carex aquatilis	3	Potentilla glandulosa	1
Carex nebrascensis	3,6	Potentilla gracilis	1,2,5
Carex praegracilis	3,6	Rosa woodsii	1,4
Carex utriculata	3,6	Rumex crispus	2,5,6
Centaurea maculosa	1	Salix bebbiana	4,6
Cirsium arvense	1,2,5	Salix exigua	4,6
Cornus stolonifera	4	Salix lemmonii	4,6
Deschampsia cespitosa		Scirpus acutus	3,6
Eleocharis palustris	2,4,5,6	Senecio sphaerocephalus	5,6
Eleocharis pauciflora	5,6	Sisyrinchium angustifolium	5
Epilobium ciliatum	2,5,6	Taraxacum officinale	1
Equisetum arvense	1,2,5	Thlaspi arvense	1
Festuca octoflora	1	Trifolium pratense	1
Geum Aleppicum	2,5,6	Triglochin palustre	3,5,6
Glyceria elata	_	Typha latifolia	3,6
Gnaphalium palustre	2,5		
Hippuris vulgaris	2,6		
Hordeum brachyantherum	2,5,6		
Hordeum jubatum	1,2,5		

Comments / Problems: _____

PLANTED WOODY VEGETATION SURVIVAL

Plant Species	Number Originally Planted	Number Observed	Mortality Causes
Thin-leaf Alder	470	261	Doing ok
Water Birch	245	4	nearly 100% mortality
Red-oiser Dogwood	246	73	30% survival.

Comments / Problems: Greater than 50% of the alder survived through year 2 but the dogwood and birch have not fared well. Some planting locations appeared to be too wet and the birch and dogwood were also outcompeted by herbaceous forbs and grasses. Birch and dogwood are not common in the project area and may not be suited for that particular environment.

WILDLIFE

Birds

Were man-made nesting structures installed? Yes

If yes, type of structure: **blue bird and wood duck boxes** How many? **14 total**

Are the nesting structures being used? <u>Yes</u> Do the nesting structures need repairs? **No**

Mammals and Herptiles

Mammal and Harntila Species	Number	umber Indirect Indication of Use				
Mammal and Herptile Species	Observed	Tracks	Scat	Burrows	Other	
deer	0	\boxtimes	\boxtimes			
moose	3		\boxtimes			
badger	0			\boxtimes		

Additional Activities Checklist:

NA Macroinvertebrate Sampling (if required)

Comments / Problems: Blue bird and wood duck boxes installed late summer 2008 - most were unused in 2009, except two blue bird boxes that according to the landowner were used by tree swallows. Boxes were all in good condition. A cow and calf moose were seen onsite along with a single cow moose. Landowner documented a northern harrier nest in the northwest portion of the mitigation site - adults and young were seen foraging within project boundaries during the monitoring.

PHOTOGRAPHS

Using a camera with a 50mm lens and color film take photographs of the following permanent reference points listed in the check list below. Record the direction of the photograph using a compass. When at the site for the first time, establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3 feet above ground. Survey the location with a resource grade GPS and mark the location on the aerial photograph.

Photograph Checklist:

\geq		One	photogra	aph for	r each	of th	e four	cardinal	directions	surroun	ding	the wetl	land.	
	7													

At least one photograph showing upland use surrounding the wetland. If more than one upland exists then take additional photographs.

At least one photograph showing the buffer surrounding the wetland.

One photograph from each end of the vegetation transect, showing the transect.

Location	Photograph Frame #	Photograph Description	Compass Reading (°)
PP1		View looking north	345
PP1		View looking west	270
PP1		View looking east	80
PP2		View looking northwest	300
PP2		View looking west	260
PP2		View looking east	80
PP3		View looking southeast	165
PP3		View looking west	270
PP3		View looking northwest	320
PP4		View looking southeast	165
PP4		View looking southeast	100
PP4		View looking southwest	210
PP5		View looking east	90
PP5		View looking southwest	245
PP5		View looking northwest	310
PP6		View looking southeast	110
PP6		View looking south	180
PP6		View looking northwest	320
PP7		View looking southwest	200
PP7		View looking west	270
PP7		View looking south	180
Transect 1		View from start of Transect looking north	5
Transect 1		View from end of Transect looking south	185
North End		View of woody plantings	

Comments	/	Prob.	lems:	

GPS SURVEYING

Using a resource grade GPS survey the items on the checklist below. Collect at least 3 location points set at a 5 second recording rate. Record file numbers for site in designated GPS field notebook.

GPS Checklist:
☐ Jurisdictional wetland boundary.
4-6 landmarks that are recognizable on the aerial photograph.
Start and End points of vegetation transect(s).
Photograph reference points.
Groundwater monitoring well locations.
Comments / Problems:
WETLAND DELINEATION
(attach COE delineation forms)
At each site conduct these checklist items:
Delineate wetlands according to the 1987 Army COE manual.
Delineate wetland – upland boundary onto aerial photograph.
Yes Survey wetland – upland boundary with a resource grade GPS survey.
Comments / Problems:
FUNCTIONAL ASSESSMENT
(Complete and attach full MDT Montana Wetland Assessment Method field forms.)
(Also attach any completed abbreviated field forms, if used)
(This allacit any completed abore vialed forms, if asea)
Comments / Problems:
MAINTENANCE
Were man-made nesting structure installed at this site? Yes
If yes, do they need to be repaired? No
If yes, describe the problems below and indicate if any actions were taken to remedy the problems.
Were man-made structures built or installed to impound water or control water flow into or out of the
wetland? <u>No</u>
If yes, are the structures working properly and in good working order? <u>NA</u>
If no, describe the problems below.
Comments / Problems: Nesting structures installed after monitoring in 2008 (MDT communication).

MDT WETLAND MONITORING - VEGETATION TRANSECT

Site: <u>Big Hole Grazing Association</u> Date: <u>August 12, 2009</u> Examiner: <u>PBS&J - Traxler</u>
Transect Number: <u>1</u> Approximate Transect Length: <u>1247 feet</u> Compass Direction from Start: <u>5</u> Note: <u>Page 1</u>

Vegetation Type: Type 1 Upland					
Length of transect in this type: 45 feet					
Plant Species		Cover			
Agrostis alba		2 = 6-10%			
Festuca octoflora		3 = 11-20%			
Poa pratensis		3 = 11-20%			
Phleum pratense		3 = 11-20%			
Taraxacum officinale		1 = 1-5%			
Tota	al Vegetative Cover:	90%			

Vegetation Type: Type 5 Wet Meadow	
Length of transect in this type: 30 feet	
Plant Species	Cover
Agrostis alba	2 = 6-10%
Hordeum brachyantherum	2 = 6-10%
Glyceria elata	2 = 6-10%
Alopecurus pratensis	4 = 21-50%
Calamagrostis canadensis	1 = 1-5%
Senecio sphaerocephalus	1 = 1-5%
Myosotis scorpioides	2 = 6-10%
Total Vegetative Cover:	80%

Vegetation Type: Type 1 Upland	
Length of transect in this type: 230 feet	
Plant Species	Cover
Poa juncifolia	2 = 6-10%
Agropyron trachycaulum	1 = 1-5%
Hordeum jubatum	2 = 6-10%
Achillea millefolium	1 = 1-5%
Phleum pratense	3 = 11-20%
Alopecurus pratensis	1 = 1-5%
Total Vegetative Cover:	75%

Vegetation Type: Type 5 Wet Meadow	
Length of transect in this type: 320 feet	
Plant Species	Cover
Agrostis alba	3 = 11-20%
Alopecurus pratensis	1 = 1-5%
Glyceria elata	2 = 6-10%
Senecio sphaerocephalus	1 = 1-5%
Hordeum brachyantherum	2 = 6-10%
Juneus balticus	3 = 11-20%
Carex aqautilis	2 = 6-10%
Total Vegetative Cover:	75%

MDT WETLAND MONITORING - VEGETATION TRANSECT

Site: <u>Big Hole Grazing Association</u> Date: <u>August 12, 2009</u> Examiner: <u>PBS&J - Traxler</u>
Transect Number: <u>1</u> Approximate Transect Length: <u>1247 feet</u> Compass Direction from Start: <u>5</u> Note: <u>Page 2</u>

Vegetation Type: Type 1 Upland	
Length of transect in this type: 240 feet	
Plant Species	Cover
Agropyron trachycaulum	2 = 6-10%
Phleum pratense	3 = 11-20%
Taraxacum officinale	2 = 6-10%
Potentilla gracilis	2 = 6-10%
Agrostis alba	3 = 11-20%
Juncus balticus	2 = 6-10%
Alopecurus pratensis	2 = 6-10%
Total Vegetative Cover:	75%

Vegetation Type: Type 5 Wet Meadow	
Length of transect in this type: 181 feet	
Plant Species	Cover
Agrostis alba	2 = 6-10%
Glyceria elata	3 = 11-20%
Alopecurus pratensis	3 = 11-20%
Hordeum brachyantherum	2 = 6-10%
Epilobium ciliatum	1 = 1-5%
Deschampsia cespitosa	1 = 1-5%
Beckmannia syzigachne	2 = 6-10%
Senecio sphaerocephalus	2 = 6-10%
Total Vegetative Cover:	75%

Vegetation Type: Type 1 Upland	
Length of transect in this type: 51 feet	
Plant Species	Cover
Agrostis alba	3 = 11-20%
Poa juncifolia	3 = 11-20%
Agropyron trachycaulum	1 = 1-5%
Phleum pratense	2 = 6-10%
Taraxacum officinale	1 = 1-5%
Hordeum jubatum	2 = 6-10%
Total Vegetative Cover:	80%

Vegetation Type: Type 3/5 Wetland			
Length of transect in this type: 150 feet			
Plant Species	Cover		
Agrostis alba	1 = 1-5%		
Epilobium ciliatum	1 = 1-5%		
Rumex crispus	+=<1%		
Deschampsia cespitosa	2 = 6-10%		
Carex nebrascensis	1 = 1-5%		
Beckmannia syzigachne	2 = 6-10%		
Total Vegetative Cover:	75%		

MDT WETLAND MONITORING - VEGETATION TRANSECT

Cover Estima	te	Indicator Class	Source
+=<1%	3 = 11-10%	+ = Obligate	P = Planted
1 = 1-5%	4 = 21-50%	- = Facultative/Wet	V = Volunteer
2 - 6 - 10%	5 -> 50%	0 - Facultative	

Percent of perimeter developing wetland vegetation (excluding dam/berm structures): 70%

Establish transects perpendicular to the shoreline (or saturated perimeter). The transect should begin in the upland area. Permanently mark this location with a standard metal fencepost. Extend the imaginary transect line towards the center of the wetland, ending at the 3 foot depth (in open water), or at the point where water depths or saturation are maximized. Mark this location with another metal fencepost.

Estimate cover within a 10 foot wide "belt" along the transect length. At a minimum, establish a transect at the windward and leeward sides of the wetland. Remember that the purpose of this sampling is to monitor, not inventory, representative portions of the wetland site.

Comments:	
-----------	--

BIRD SURVEY - FIELD DATA SHEET

Site: **Big Hole Grazing Association** Date: **8/12/09**

Survey Time: <u>7:30</u> am to <u>1:00</u> pm

Bird Species	#	Behavior	Habitat	Bird Species	#	Behavior	Habitat
Mallard	5	L	MA				
Northern Harrier	4	FN	MA UP				
Common Snipe	3	FL	MA				
Sandhill Crane	4	FO					
Song Sparrow	2	F	UP SS				
Tree Swallow	12	F	WM UP				
Red-winged Blackbird	6	FN	MA SS				
Magpie	1	L	SS				
Raven	1	FO					
Rock Dove	3	FO					

BEHAVIOR CODES

BP = One of a breeding pair **BD** = Breeding display

 $\mathbf{F} =$ Foraging

FO = Flyover **L** = Loafing

N = Nesting

HABITAT CODES

AB = Aquatic bed
FO = Forested
I = Island
WM = Wet meadow
WA = Marsh
US = Unconsolidated shore

MF = Mud Flat OW = Open Water

Weather: Partly Sunny and warm

Notes: Land Owner documented Northern Harrier nest in NW portion of mitigation site.

DATA FORM ROUTINE WETLAND DETERMINATION

(1987 COE Wetlands Delineation Manual)

Project/Site: Big Hole Grazing Association

Applicant/Owner: Montana Department of Transportation

Investigators: PBSJ - Traxler

Project No:

Date: 12-Aug-2009 County: Beaverhead State: Montana

State: Mon Plot ID: 1

Do Normal Circumstances exist on the site?

Is the site significantly disturbed (Atypical Situation:)?

Is the area a potential Problem Area?
(If needed, explain on the reverse side)

Yes No Yes No Yes No Community ID: EM/SS

Transect ID: Field Location: Rock Creek drainage

VEGETATION

(USFWS Region No. 9)

YEOLIATION			igion 140. 3)		
Dominant Plant Species(Latin/Common)	Stratum	Indicator	Plant Species(Latin/Common)	Stratum	Indicator
Salix bebbiana	Shrub	FACW	Agrostis alba	Herb	FACW
Willow,Bebb			Redtop		
Salix exigua	Shrub	OBL	Juncus balticus	Herb	OBL
Willow,Sandbar			Rush,Baltic		
Salix lemmonii	Shrub	FACW+	Carex aquatilis	Herb	OBL
Willow,Lemmon's	7		Sedge,Water		1
Alnus incana	Shrub	FACW	Calamagrostis canadensis	Herb	FACW+
Alder,Speckled	7		Reedgrass,Blue-Joint		
Betula pumila	Shrub	OBL	Hippuris vulgaris	Herb	OBL
Birch,Bog			Mare's-Tail,Common		l
Comus stolonifera	Shrub	FACW			
Dogwood,Red-Osier					
	_				
		I			

Percent of Dominant Species that are OBL, FACW or FAC: (excluding FAC-) 11/11 = 100.00%

FAC Neutral: 11/11 = 100.00%

Numeric Index: 17/11 = 1.55

Remarks:

Rock Creek floodplain has significant SS component with emergent marsh understory.

HYDROLOGY

NO Recorded Data(Describe in Remarks):		Wetland Hydrology Indicators
N/A Stream, Lake or Tide Gau	ge	Primary Indicators
<u>N/A</u> Aerial Photographs		NO Inundated
<u>N/A</u> Other		YES Saturated in Upper 12 Inches
VES No Becarded Date		NO Water Marks
YES No Recorded Data		YES Drift Lines
		YES Sediment Deposits
Field Observations		YES Drainage Patterns in Wetlands
		Secondary Indicators
Depth of Surface Water:	N/A (in.)	NO Oxidized Root Channels in Upper 12 Inches
	N1/A # 1	NO Water-Stained Leaves
Depth to Free Water in Pit:	N/A (in.)	NO Local Soil Survey Data
Depth to Saturated Soil: = 6 (in.)		YES FAC-Neutral Test
Depth to Saturated Soil.	- 0 (III.)	NO Other(Explain in Remarks)

Remarks:

Plot taken 5 feet from channel. Channel water depth varies from 1-36 inches. Conditions unchanged from 2008.

Page 1 of 2 WetFormtm

DATA FORM ROUTINE WETLAND DETERMINATION

(1987 COE Wetlands Delineation Manual)

(1967 COE Wetlands Delineation Wandar)					
Project/Site: Big Hole Grazing Association Applicant/Owner: Montana Department of Transportation Investigators: PBSJ - Traxler			Project No:	Date: 12-Aug-2009 County: Beaverhead State: Montana Plot ID: 1	
SOILS					
Map Unit Name (Series and Phase):	Mooseflat loam, 0-4% slo	pe		

Depth inches)	Horizon	Matrix Color (Munsell Moist)	Mottle Color (Munsell Moist)	Mot Abundance		Texture, Concretions, Structure, etc
4	Α	10YR3/2	N/A	N/A	N/A	Loam, organics
14	В	10YR4/2	10YR5/8	Common	Distinct	Sandy loam, small cobbles
Hydric Soil Indicators: NO Histosol NO Histic Epipedon YES Sulfidic Odor NO Aquic Moisture Regime NO Reducing Conditions YES Gleyed or Low Chroma Colors			NO Concretions NO High Organic Content in Surface Layer in Sandy Soils NO Organic Streaking in Sandy Soils NO Listed on Local Hydric Soils List NO Listed on National Hydric Soils List NO Other (Explain in Remarks)			

WETLAND DETERMINATION

Hydrophytic Vegetation Present? Wetland Hydrology Present? Hydric Soils Present?	Yes No Yes No Yes No	Is the Sampling Point within the Wetland? (Yes) No
Remarks:		

Page 2 of 2 WetForm^{lm}

DATA FORM ROUTINE WETLAND DETERMINATION

(1987 COE Wetlands Delineation Manual)

Project/Site: Big Hole Grazing Association

Applicant/Owner: Montana Department of Transportation

Investigators: PBS&J - Traxler

Project No:

Date: 12-Aug-2009 County: Beaverhead

State: Montana

Plot ID: 2

Do Normal Circumstances exist on the site?

(If needed, explain on the reverse side)

is the site significantly disturbed (Atypical Situation:)?

is the area a potential Problem Area?

Yes No

(No)

Community ID: Wet Meadow

Transect ID: Field Location:

Wet meadow towards center of site

VEGETATION

(USFWS Region No. 9)

Yes

Dominant Plant Species(Latin/Common)	Stratum	Indicator	Plant Species(Latin/Common)	Stratum	Indicator
Agrostis alba	Herb	FACW	Juncus balticus	Herb	OBL
Redtop			Rush,Baltic	1	İ
Alopecurus pratensis	Herb	FACW	Juncus longistylis	Herb	FACW
Foxtail,Meadow			Rush,Long-Style	1	
Carex nebrascensis	Herb	OBL	Senecio sphaerocephalus	Herb	FACW
Sedge,Nebraska	1		Groundsel,Ball-Head	1	
Hordeum brachyantherum	Herb	FACW			
Barley,Meadow					
				_	
				4	
	1				
	4			4	
Dave and of Davis and Creation that are OBI		<u> </u>	7/7 = 100 009/		

Percent of Dominant Species that are OBL, FACW or FAC: (excluding FAC-) 7/7 = 100.00%

FAC Neutral:

7/7 = 100.00%

Numeric Index: 12/7 = 1.71

Remarks:

Flat wet meadow wetland towards center of site.

HYDROLOGY

NO Recorded Data(Describe in Rem	arks):	Wetland Hydrology Indicators	
<u>N/A</u> Stream, Lake or Tide Gaug	e	Primary Indicators	
<u>N/A</u> Aerial Photographs		NO Inundated	
<u>N/A</u> Other		YES Saturated in Upper 12 Inches	
YES No Recorded Data		NO Water Marks	1
TES NO Recorded Data		NO Drift Lines	1
		NO Sediment Deposits	
Field Observations		NO Drainage Patterns in Wetlands	
		Secondary Indicators	
Depth of Surface Water:	N/A (in.)	NO Oxidized Root Channels in Upper 12 Inches	
	NI/A a v	NO Water-Stained Leaves	
Depth to Free Water in Pit:	N/A <i>(in.)</i>	NO Local Soil Survey Data	
Depth to Saturated Soil:	= 0 (in.)	YES FAC-Neutral Test	
Depth to Saturated Soil.	- O (III.)	NO Other(Explain in Remarks)	

Remarks:

high groudwater table. Water levels increased across site with plugging of drainage ditches. Ground saturated to surface in 2009.

Page 1 of 2 WetFormtm

DATA FORM ROUTINE WETLAND DETERMINATION

(1987 COE Wetlands Delineation Manual)

Project/Site: Big Hole Grazing Association Project No: Date: 12-Aug-2009
Applicant/Owner: Montana Department of Transportation County: Beaverhead State: Montana Plot ID: 2

SOILS

Depth (inches)	· 1		Mottle Color (Munsell Moist)	Mot Abundance		Texture, Concretions, Structure, etc
6	Α	10YR3/2	N/A	N/A	N/A	Loam, Roots
12	В	10YR3/1	10YR6/8	Common	Distinct	Clay loam
	NO Sulfic NO Aquic NO Redu YES Gleye	: Epipedon		NO High NO Org NO List NO List	anic Streak ed on Loca ed on Natic	content in Surface Layer in Sandy Soils ing in Sandy Soils I Hydric Soils List onal Hydric Soils List in Remarks)
Remarks	; : ,					

Hydrophytic Vegetation Present?	(Yes) No	Is the Sampling Point within the Wetland? (Yes) No
Wetland Hydrology Present?	(Yes) No	
Hydric Soils Present?	(Yes) No	
Remarks:		

Page 2 of 2 WetForm^{im}

DATA FORM ROUTINE WETLAND DETERMINATION

(1987 COE Wetlands Delineation Manual)

Project/Site: Big Hole Grazing Association

Applicant/Owner: Montana Department of Transportation

Investigators: PBS&J - Traxler

Project No:

Date: 12-Aug-2009 County: Beaverhead

State: Montana

Plot ID: 3

Do Normal Circumstances exist on the site?

(If needed, explain on the reverse side)

Is the site significantly disturbed (Atypical Situation:)?

Is the area a potential Problem Area?

Yes Yes

(Yes)

No Co

Community ID: EM Transect ID:

Field Location:

East side of mitigation area

VEGETATION

(USFWS Region No. 9)

Dominant Plant Species(Latin/Common)	Stratum	Indicator	Plant Species(Latin/Common)	Stratum	Indicato
Carex aquatilis	Herb	OBL	Juncus balticus	Herb	OBL
Sedge,Water	1		Rush,Baltic		
Carex nebrascensis	Herb	OBL	Geum aleppicum	Herb	FACW-
Sedge,Nebraska			Avens, Yellow		
Carex rostrata	Herb	OBL	Salix bebbiana	Shrub	FACW
Sedge,Beaked			Willow,Bebb		
Agrostis alba	Herb	FACW	Salix exigua	Shrub	OBL
Redtop			Willow,Sandbar		ļ
		<u> </u>			
	-				

Percent of Dominant Species that are OBL, FACW or FAC:

FAC Neutral:

8/8 = 100.00%

(excluding FAC-) 8/8 = 100.00%

Numeric Index:

11/8 = 1.38

Remarks:

Site is converting from heavily grazed emergent marsh to EM/SS. Significant hummocks throughout.

HYDROLOGY

NO Recorded Data(Describe in Rer	narks):	Wetland Hydrology Indicators					
N/A Stream, Lake or Tide Gauge		Primary Indicators					
<u>N/A</u> Aerial Photographs		NO Inundated					
<u>N/A</u> Other		YES Saturated in Upper 12 Inches					
VEC No Broaded Date		NO Water Marks					
YES No Recorded Data		NO Drift Lines					
		NO Sediment Deposits					
Field Observations		NO Drainage Patterns in Wetlands					
		Secondary Indicators					
Depth of Surface Water:	N/A (in.)	NO Oxidized Root Channels in Upper 12 Inches					
	N1/A 22 h	NO Water-Stained Leaves					
Depth to Free Water in Pit:	N/A (in.)	NO Local Soil Survey Data					
Depth to Saturated Soil:	= 0 (in.)	YES FAC-Neutral Test					
Deptil to Saturated Soil.	- O (III.)	<u>NO</u> Other(Explain in Remarks)					

Remarks:

High groundwater table. Similar conditions to 2008.

Page 1 of 2 WetFormtm

DATA FORM ROUTINE WETLAND DETERMINATION

(1987 COE Wetlands Delineation Manual)

Project/Site: Big Hole Grazing Association Project No: Date: 12-Aug-2009 Applicant/Owner: Montana Department of Transportation County: Beaverhead Investigators: PBS&J - Traxler State: Montana Plot ID: 3 SOILS Foxgulch-copperbasin-Wisdom complex Map Unit Name (Series and Phase): Map Symbol: 15A Drainage Class: Somewhat poorly drained Mapped Hydric Inclusion? Taxonomy (Subgroup): Field Observations Confirm Mapped Type? Yes No. **Profile Description** Depth Matrix Color **Mottle Color** Mottle (inches) Horizon Texture, Concretions, Structure, etc (Munsell Moist) (Munsell Moist) Abundance/Contrast 14 A/B 10YR2/1 N/A Clay loam Hydric Soil Indicators: NO Histosol **NO Concretions** NO Histic Epipedon NO High Organic Content in Surface Layer in Sandy Soils NO Sulfidic Odor NO Organic Streaking in Sandy Soils

NO Listed on Local Hydric Soils List

NO Other (Explain in Remarks)

NO Listed on National Hydric Soils List

Re	m	ar	ks	:

WETLAND DETERMINATION

NO Aquic Moisture Regime

YES Gleyed or Low Chroma Colors

NO Reducing Conditions

Hydrophytic Vegetation Present? Wetland Hydrology Present? Hydric Soils Present?	Yes Yes Yes	No No No	Is the Sampling Point within the Wetland?	(es)	No
Remarks:					

Page 2 of 2 WetFormtm

MDT Montana Wetland Assessment Form (revised March 2008)

1.	Project Name: Big Hole Gra	azing Association 2. MDT Pro	oject #: <u>STPX 1(45)</u> 3. Contr	ol #: <u>CN4668</u>									
3.	Evaluation Date: 8/12/09 4. Evaluator(s): PBS&J - Traxler 5. Wetland/Site #(s): Rock Creek corridor												
6.	Wetland Location(s): Township 4 S, Range 16 W, Section 2; Township N, Range E, Section												
	Approximate Stationing or Roadposts:												
	Watershed: 6 - Upper Miss	souri County: Beaverhead	·										
7.	Evaluating Agency: MDT Purpose of Evaluation: (measured, e.g. GPS) Wetland potentially affected by MDT project Mitigation wetlands; pre-construction Mitigation wetlands; post-construction Other (see manual for determining AA) (measured, e.g. GPS)												
	Mitigation wetlands;												
10		oost-construction TLAND AND AQUATIC HABI	(see manual for de TATS IN AA (See manual for d	etermining AA) (mea lefinitions.)	sured, e.g. GPS)								
10	☑ Mitigation wetlands; p☐ Other	oost-construction	(see manual for de	etermining AA) (mea									
10		oost-construction TLAND AND AQUATIC HABI	(see manual for de TATS IN AA (See manual for d	etermining AA) (mea lefinitions.)	sured, e.g. GPS)								
10		oost-construction TLAND AND AQUATIC HABI Class (Cowardin)	(see manual for de TATS IN AA (See manual for d	termining AA) (mea lefinitions.) Water Regime	sured, e.g. GPS)								
10	Mitigation wetlands; p Other CLASSIFICATION OF WE HGM Class (Brinson) Riverine	TLAND AND AQUATIC HABIT Class (Cowardin) Rock Bottom	(see manual for de TATS IN AA (See manual for d	termining AA) (mea lefinitions.) Water Regime Permanent / Perennial	sured, e.g. GPS) % OF AA 10								
10	Mitigation wetlands; p Other CLASSIFICATION OF WE HGM Class (Brinson) Riverine Riverine	TLAND AND AQUATIC HABIT Class (Cowardin) Rock Bottom Emergent Wetland	(see manual for de TATS IN AA (See manual for d	termining AA) (mean lefinitions.) Water Regime Permanent / Perennial Seasonal / Intermittent	% OF AA 10 20								
10	Mitigation wetlands; p Other CLASSIFICATION OF WE HGM Class (Brinson) Riverine Riverine	TLAND AND AQUATIC HABIT Class (Cowardin) Rock Bottom Emergent Wetland	(see manual for de TATS IN AA (See manual for d	termining AA) (mean lefinitions.) Water Regime Permanent / Perennial Seasonal / Intermittent	% OF AA 10 20								
10	Mitigation wetlands; p Other CLASSIFICATION OF WE HGM Class (Brinson) Riverine Riverine	TLAND AND AQUATIC HABIT Class (Cowardin) Rock Bottom Emergent Wetland	(see manual for de TATS IN AA (See manual for d	termining AA) (mean lefinitions.) Water Regime Permanent / Perennial Seasonal / Intermittent	% OF AA 10 20								

Comments: ____

11. ESTIMATED RELATIVE ABUNDANCE (of similarly classified sites within the same Major Montana Watershed Basin; see manual.) common

12. GENERAL CONDITION OF AA

i. Disturbance: Use matrix below to select the appropriate response; see manual for Montana listed noxious weed and aquatic nuisance vegetation species lists.

	Predominant Conditions Adjacent to (within 500 feet of) AA								
Conditions within AA	Managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or buildings; and noxious weed or ANVS cover is ≤15%.	Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >30%.							
AA occurs and is managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings; and noxious weed or ANVS cover is ≤15%.		low disturbance							
AA not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to relatively minor clearing, fill placement, or hydrological alteration; contains few roads or buildings; noxious weed or ANVS cover is ≤30%.									
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >30%.									

Comments (types of disturbance, intensity, season, etc.): Area is currently fenced and not subject to grazing or other forms of disturbance.

ii. Prominent noxious, aquatic nuisance, and other exotic vegetation species:

13. STRUCTURAL DIVERSITY (Based on number of "Cowardin" vegetated classes present [do not include unvegetated classes]; see #10 above.)

Existing # of "Cowardin" Vegetated Classes in AA	Initial Rating	Is current management peristence of additional		Modified Rating
≥3 (or 2 if one is forested) classes		NA	NA	NA
2 (or 1 if forested) classes	mod	NA	NA	NA
1 class, but not a monoculture		←NO	YES→	
1 class, monoculture (1 species comprises ≥90% of total cover)		NA	NA	NA

Comments: Two vegetated classes in creek corridor.

iii. Provide brief descriptive summary of AA and surrounding land use/habitat: AA includes the Rock Creek channel and adjacent SS and EM wetland. Surrounding land outside project area is grazed and hayed.

14A. HABITAT FOR FEDER	ALLY	LISTE	D OR	PRO	POSE	D THE	REATE	NED	OR E	NDAN	GERE	D PL	ANTS	OR A	NIMAL	.s				
. AA is Documented (D) or Suspected (S) to contain: Check box based on definitions in manual. Primary or critical habitat (list species)																				
ii. Rating: Based on the strong	ngest h	abitat	chose	n in	14A(i) a	above,	, selec	t the	corres	pondin	ig func	tiona	l point	and ra	ting.					
Highest Habitat Level	Doc/F	Primai	ry S	us/P	rimary	Do	c/Sec	onda	ry S	us/Se	conda	ry	Doc/In	ciden	tal	Sus/	Incide	ntal	None	•
Functional Point/Rating				-						-									0L	
Sources for documented us	Sources for documented use (e.g. observations, records): <u>.</u>																			
14B. HABITAT FOR PLANTS OR ANIMALS RATED S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM Do not include species listed in 14A above.																				
i. AA is Documented (D) or Suspected (S) to contain: Check box based on definitions in manual. Primary or critical habitat (list species)																				
ii. Rating: Based on the stro	ngest	habita	t chose	en in	14A(i)	above	, selec	ct the	corres	spondii	ng fund	ctiona	al point	and ra	ating.					_
Highest Habitat Level	Doc/F	Primai	ry S	us/P	rimary	Do	c/Sec	onda	ry S	us/Se	conda	ry	Doc/In	ciden	tal S	Sus/I	ncider	ntal	None	_
S1 Species Functional Point/Rating	•			-						-				.2L						
S2 and S3 Species Functional Point/Rating				-						-										
Sources for documented us	e (e n	ohsei	rvation	s red	ords).	MEW	P - MF	ISH												J
i. Evidence of Overall Wildl Substantial: Based on an observations of abunda abundant wildlife sign sopresence of extremely interview with local biol	y of the ant wild such as limiting ogist w	e in the followallife #s scat, habita	wing [c or hig tracks at feat owledo	check gh spe s, nes ures ge of]. ecies d t struct not ava	iversit ures, ailable	y (duri	ng ar trails,	ny perio	od)	M	inima few little spar	orting e al: Bas or no v to no v se adja view w	ed on vildlife vildlife acent u	any of observ sign upland	vatior food	source	ng pea es	k use	
 Moderate: Based on any of sobservations of scatter Scommon occurrence of adequate adjacent uplar interview with local biol Wildlife Habitat Features For class cover to be conside percent composition of the AAS/I = seasonal/intermittent; T/ 	ed wild wildlife and foo ogist w : Work red eve A (see #	life gro e sign d sour ith kno ing fro enly di #10).	oups o such a rces owledg om top stribut Abbre	r indicas sca ge of to bo ed, th viatio	at, track the AA ottom, cone mosins for s	cs, nes heck a t and l surface	appropleast per water	oriate reval er dura	AA attent verations	e trails tributes getate are as	s in ma d class follow	trix to ses m s: P/F	o arrive nust be	within maner	20% ont/pere	of ead	ch othe			
Structural Diversity		прога	у/српс			Λ = α	DOCITE	[3CC	Tiariua	110114					terrinoj	1				
(see #13)				الا	High				ļ		Ľ	7 INIO	derate						OW	
Class Cover Distribution (all vegetated classes)			ven			☐ Un	even			□ E	ven			🛛 Un	even			□ E	ven	
Duration of Surface Water in ≥ 10% of AA	P/P	S/I	T/E	Α	P/P	S/I	T/E	Α	P/P	S/I	T/E	Α	P/P	S/I	T/E	Α	P/P	S/I	T/E	A
□ Low Disturbance at AA	i							l					Е							
(see #12i)													_							
☐ Moderate Disturbance at AA (see #12i)																				
☐ High Disturbance at AA (see #12i)																				
iii. Rating: Use the conclusi	ons fro	m i an	ıd ii ab	ove a	and the	matri	x belov	w to s	select t	he fun	ctional	poin	t and ra	ating.						
Evidence of Wildlife Use										s Ratir										
(i)		⊠ Exc	ceptio	nal			High				derate	•		☐ Lo	W	_				
Substantial										-						1				
Moderate			.9H													-				
Minimal		- U -	 - al I			L				-						╛				
Comments: various bird spec	cies, sr	nail ar	nd larg	e ma	mmals.	nerps	<u>S.</u>													

14D. GENERAL FISH HAE	BITAT 🗌 NA	(proceed to 14E)
-----------------------	------------	------------------

If the AA is not used by fish, fish use is not restorable due to habitat constraints, or is not desired from a management perspective [such as fish entrapped in a canal], then check the NA box and proceed to 14E.

Assess this function if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [i.e., fish use is precluded by perched culvert or other barrier].

Type of Fishery: Cold Water (CW) Warm Water (WW) Use the CW or WW guidelines in the manual to complete the matrix.

i. Habitat Quality and Known / Suspected Fish Species in AA: Use matrix to select the functional point and rating.

Duration of Surface Water in AA	⊠ Pe	erman	ent / P	erenn	ial		□s	easor	nal / Ir	ntermit	tent	•	П.	empo	rary / E	Ephen	neral	
Aquatic Hiding / Resting / Escape Cover	Opti] imal	∑ Adeq	uate	Po	oor	Opti] imal	Ade] quate	Po	or	Op	 timal	Aded] uate	Po	oor
Thermal Cover: optimal / suboptimal	0	S	0	S	0	S	0	S	0	S	0	S	0	S	0	s	0	S
FWP Tier I fish species			.8H															
FWP Tier II or Native Game fish species																		
FWP Tier III or Introduced Game fish																		
FWP Non-Game Tier IV or No fish species																		

Sources used for identifying fish spp. potentially found in AA: 2008 MFISH on FWP website. Westslope cutthroat, grayling, brook trout, brown trout, rainbow trout, whitefish.

ii.	Modified Rating:	NOTE: Modified score cannot exceed	1.0 or be less than 0.1
-----	------------------	------------------------------------	-------------------------

a) Is fish use of the AA significantly reduced by a culvert, dike, or other man-made structure or activity, or is the waterbody included on the current final MDEQ list of waterbodies in need of TMDL development with listed "Probable Impaired Uses" including cold or warm water fishery or aquatic life support, or do aquatic nuisance plant or animal species (see Appendix E) occur in fish habitat? TyES, reduce score in i by 0.1 = or 🖾 N0

b) Does the AA contain a documented spawning area or other critical habitat feature (i.e., sanctuary pool, upwelling area; specify in comments) for native fish or introduced game fish? \square YES, add to score in i or iia 0.1 = or \boxtimes N0

iii.	Final	Score	and	Rating:	.8H	Comments:	
------	--------------	--------------	-----	---------	-----	-----------	--

14E. FLOOD ATTENUATION


□ NA (proceed to 14F)


Applies only to wetlands that are subject to flooding via in-channel or overbank flow. If wetlands in AA are not flooded from in-channel or overbank flow, check the NA box and proceed to 14F.

Entrenchment Ratio (ER) Estimation (see manual for additional guidance). Entrenchment ratio = (flood-prone width) / (bankfull width).

Flood-prone width = estimated horizontal projection of where 2 X maximum bankfull depth elevation intersects the floodplain on each side of the stream.

flood prone width / bankfull width = entrenchment ratio





i. Rating: Working from top to bottom, use the matrix below to select the functional point and ratina.

Estimated or Calculated Entrenchment					erately Entr				
(Rosgen 1994, 1996)	C, D, E stream types		В	stream typ	е	A, F, G stream types			
Percent of Flooded Wetland Classified as									
Forested and/or Scrub/Shrub	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
AA contains no outlet or restricted outlet									
AA contains unrestricted outlet		.8H							

ii. Are ≥10 acres of wetland in the AA subject to flooding AND are man-made features which may be significantly damaged by floods located within 0.5 mile downstream of the AA?

YES

NO Comments: no structures

14F.	SHORT AND LONG TERM SURFACE WATER STORAGE	☐ NA (proceed to 14G)
	Applies to wetlands that flood or pond from overbank or in-chann	nel flow, precipitation, upland surface flow, or groundwater flow.
	If no wetlands in the AA are subject to flooding or ponding, then	check the NA box and proceed to 14G.

i. Rating: Working from top to bottom, use the matrix below to select the functional point and rating. Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; and T/E = temporary/ephemeral [see manual for further definitions of these terms].

Estimated Maximum Acre Feet of Water Contained in Wetlands within the AA that are Subject to Periodic Flooding or Ponding	☐ >5 acre feet		eet	☑ 1.1 to 5 acre feet			☐ ≤1 acre foot		
Duration of Surface Water at Wetlands within the AA	□ P/P	□ S/I	□ T/E	⊠ P/P	□ S/I	□ T/E	□ P/P	□ S/I	□ T/E
Wetlands in AA flood or pond ≥ 5 out of 10 years				.8H					
Wetlands in AA flood or pond < 5 out of 10 years									

^			4	۱.	_
u	mc	me	m	IS	:

14G. SEDIMENT / NUTRIENT / TOXICANT / RETENTION AND REMOV	AL NA (proceed to 14H
---	-----------------------

Applies to wetland with potential to receive sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, check the NA box and proceed to 14H.

i. Rating: Working from top to bottom, use the matrix below to select the functional point and rating.

Sediment, Nutrient, and Toxicant Input Levels within AA	AA receive has potent nutrients, such that c substantia sedimenta toxicants, present.	ial to delivor compou other funct lly impaire tion, source	er sedime nds at lev ions are n d. Minor es of nutr	ents, rels not rients or	Waterbody is on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.					
% Cover of Wetland Vegetation in AA	⊠≥∶	70%	□<	70%	□ ≥ 70%		□ < 70 %			
Evidence of Flooding / Ponding in AA	⊠ Yes	☐ No	☐ Yes	☐ No	☐ Yes	☐ No	☐ Yes	☐ No		
AA contains no or restricted outlet										
AA contains unrestricted outlet	.9H									

	ents	

14H. SEDIMENT / SHORELINE STABILIZATION NA (proceed to 14l)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body which is subject to wave action.

If 14H does not apply, check the NA box and proceed to 14I.

% Cover of Wetland Streambank or Shoreline by Species with Stability	Duration of S	ted Vegetation	
Ratings of ≥6 (see Appendix F).	Permanent / Perennial	☐ Seasonal / Intermittent	☐ Temporary / Ephemeral
⊠ ≥ 65%	1H		
□ 35-64%			
☐ < 35%			

Comments:

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. Level of Biological Activity: Synthesis of wildlife and fish habitat rates (select).

General Fish Habitat Rating	General Wildlife Habitat Rating (14Ciii)						
(14Diii)	⊠ E/H		□ L				
⊠ E/H	Н	Н					
							
□ NA							

ii. Rating: Working from top to bottom, use the matrix below to select the functional point and rating. Factor A = acreage of vegetated wetland component in the AA; Factor B = level of biological activity rating from above (14li); Factor C = whether or not the AA contains a surface or subsurface outlet; the final three rows pertain to the duration of surface water in the AA, where P/P, S/I, and T/E were previously defined, and A = "absent" [see manual for further definitions of these terms].

Α	\boxtimes	Vegeta	ted Co	mponent	>5 ac	res		Vegeta	ated Co	mponent	1-5 ac	res		Veget	ated Co	mponen	t <1 acı	re
В	⊠⊦	ligh	M	oderate		Low	_ _	ligh		derate		Low	- ⊢	ligh	☐ Mo	derate		.ow
С	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
P/P	1H																	
S/I																		
T/E/A																		

■ None

Wetland/Site #(s): Rock Creek Corridor 14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT (continued) iii. Modified Rating: Note: Modified score cannot exceed 1.0 or be less than 0.1. Vegetated Upland Buffer: Area with ≥ 30% plant cover, ≤ 15% noxious weed or ANVS cover, AND that is not subjected to periodic mechanical mowing or clearing (unless for weed control). Is there an average ≥ 50-foot wide vegetated upland buffer around ≥ 75% of the AA's perimeter? X YES, add 0.1 to score in ii = ____ NO iv. Final Score and Rating: 1H Comments: 14J. GROUNDWATER DISCHARGE / RECHARGE Check the appropriate indicators in i and ii below. i. Discharge Indicators ii. Recharge Indicators Permeable substrate present without underlying impeding layer.

Wetland contains inlet but no outlet. ☐ The AA is a slope wetland. Springs or seeps are known or observed. ☐ Vegetation growing during dormant season/drought. ☐ Stream is a known 'losing' stream. Discharge volume decreases. Wetland occurs at the toe of a natural slope. Other: ☐ Seeps are present at the wetland edge. ☐ AA permanently flooded during drought periods. Wetland contains an outlet, but no inlet. ☐ Shallow water table and the site is saturated to the surface. Other: iii. Rating: Use the information from i and ii above and the table below to select the functional point and rating Duration of Saturation at AA Wetlands FROM GROUNDWATER DISCHARGE or

WITH WATER THAT IS RECHARGING THE GROUNDWATER SYSTEM

□ T

14K.	UNIQUENESS

☐ Groundwater Discharge or Recharge

☐ Insufficient Data/Information

Criteria

Comments:

Comments:

i Rating: Working from top to bottom use the matrix below to select the functional point and rating

⊠ P/P

1H

1. Rating: Working from top to bottom, use the matrix below to select the functional point and rating.									
Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland OR plant association listed as "S1" by the MTNHP			cited ra diversi contair	es not contain are types ANI ty (#13) is hig ns plant asso as "S2" by the	Structural h OR ciation	AA does not contain previously cited rare types OR associations AND structural diversity (#13) is low-moderate		
Estimated Relative Abundance (#11)	□ Rare	☐ Common	□ Abundant	□ Rare	☐ Common	□ Abundant	□ Rare	□ Common	□ Abundant
Low Disturbance at AA (#12i)								.4M	
☐ Moderate Disturbance at AA (#12i)									
☐ High Disturbance at AA (#12i)									

☐ S/I

14L.	RECREATION / EDUCATION POTENTIAL	☐ NA (proceed to Overall Summary and Rating page)
	Affords 'bonus' points if AA provides a recreational or e	educational opportunity.

i. Is the AA a known or potential recreational or educational site? 🛛 YES, go to ii. 🗌 NO, check the NA box.

ii. Check categories that apply to the AA: ☐ Educational/Scientific Study ☐ Consumptive Recreational ☐ Non-consumptive recreational ☐ Other:

iii. Rating: Use the matrix below to select the functional point and rating.

Known or Potential Recreational or Educational Area	Known	Potential
Public ownership or public easement with general public access (no permission required)		
Private ownership with general public access (no permission required)		
Private or public ownership without general public access, or requiring permission for public access		.05L

Comments:

15. GENERAL SITE NOTES: fishing and hunting by permission.

Function & Value Variables	Rating – Actual Functional Points	Possible Functional Points	Functional Units: Actual Points x Estimated AA Acreage	Indicate the Four Most Prominent Functions with an Asterisk
A. Listed / Proposed T&E Species Habitat	low 0.00	1.00	0	
B. MT Natural Heritage Program Species Habitat	low 0.20	1.00	2	
C. General Wildlife Habitat	high 0.90	1.00	9	
D. General Fish Habitat	high 0.80	1.00	8	
E. Flood Attenuation	high 0.80	1.00	8	
F. Short and Long Term Surface Water Storage	high 0.80	1.00	8	
G. Sediment / Nutrient / Toxicant Removal	high 0.90	1.00	9	
H. Sediment / Shoreline Stabilization	high 1.00	1.00	10	
I. Production Export / Food Chain Support	high 1.00	1.00	10	
J. Groundwater Discharge / Recharge	high 1.00	1.00	10	
K. Uniqueness	mod 0.40	1.00	4	
L. Recreation / Education Potential (bonus point)	low 0.05		0.5	
Total Points	7.85	11	78.5 Total	Functional Units
Percent of Possibl	e Score 71% (round	to nearest whol	e number)	

Category I Wetland: (must satisfy one of the following criteria; otherwise go to Category II) ☐ Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; or ☐ Score of 1 functional point for Uniqueness; or ☐ Score of 1 functional point for Flood Attenuation and answer to Question 14E.ii is "yes"; or ☐ Percent of possible score > 80% (round to nearest whole #).
Category II Wetland: (Criteria for Category I not satisfied and meets any one of the following criteria; otherwise go to Category IV) ☐ Score of 1 functional point for MT Natural Heritage Program Species Habitat; or ☐ Score of .9 or 1 functional point for General Wildlife Habitat; or ☐ Score of .9 or 1 functional point for General Fish Habitat; or ☐ "High" to "Exceptional" ratings for both General Wildlife Habitat and General Fish/Aquatic Habitat; or ☐ Score of .9 functional point for Uniqueness; or ☐ Percent of possible score > 65% (round to nearest whole #).
☐ Category III Wetland: (Criteria for Categories I, II, or IV not satisfied)
Category IV Wetland: (Criteria for Categories I or II are not satisfied and all of the following criteria are met; if not go to Category III) "Low" rating for Uniqueness; and Vegetated wetland component < 1 acre (do not include upland vegetated buffer); and Percent of possible score < 35% (round to nearest whole #).
OVERALL ANALYSIS AREA (AA) RATING: Check the appropriate category based on the criteria outlined above.

MDT MONTANA WETLAND ASSESSMENT FORM (revised March 2008)

1. Project Name: Big Hole Grazing	g Association 2. MDT Pro	ject #: STPX 1(45) 3. Contro	ol #: <u>CN4668</u>							
Evaluation Date: 8/12/09 4. Evaluator(s): PBS&J - Traxler 5. Wetland/Site #(s): All wetlands outside Rock Creek corridor										
6. Wetland Location(s): Townshi	Wetland Location(s): Township 4 S, Range 16 W, Section 2; Township N, Range E, Section									
Approximate Stationing or Ro	adposts:									
Watershed: 6 - Upper Missour	i County: Beaverhead	. 								
7. Evaluating Agency: MDT Purpose of Evaluation: ☐ Wetland potentially affect ☐ Mitigation wetlands; pre-c ☑ Mitigation wetlands; post ☐ Other	construction	9. Assessment Area	e): (visually estimated) 42 (measured, e.g. GPS) (AA) Size (acre): (visuse termining AA) 39.81 (measured)							
10. CLASSIFICATION OF WETLA	AND AND AQUATIC HABIT	TATS IN AA (See manual for d	efinitions.)							
HGM Class (Brinson)	Class (Cowardin)	Modifier (Cowardin)	Water Regime	% OF AA						
Slope	Emergent Wetland		Permanent / Perennial	20						
Slope	Scrub-Shrub Wetland		Seasonal / Intermittent	10						
Slope	Emergent Wetland		Seasonal / Intermittent	70						

Comments: Springs from north and high groundwater table supply hydrology to the site

11. ESTIMATED RELATIVE ABUNDANCE (of similarly classified sites within the same Major Montana Watershed Basin; see manual.) common

12. GENERAL CONDITION OF AA

i. Disturbance: Use matrix below to select the appropriate response; see manual for Montana listed noxious weed and aquatic nuisance vegetation species lists.

	Predominant Conditions Adjacent to (within 500 feet of) AA					
Conditions within AA	state; is not grazed, nayed, logged, or otherwise converted; does not contain roads or buildings; and noxious weed or ANVS cover is \$10%.		Land cultivated or heavily grazed or logged; subject to substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANV cover is >30%.			
AA occurs and is managed in predominantly natural state; is not grazed, hayed, logged, or otherwise converted; does not contain roads or occupied buildings; and noxious weed or ANVS cover is ≤15%.		low disturbance				
AA not cultivated, but may be moderately grazed or hayed or selectively logged; or has been subject to relatively minor clearing, fill placement, or hydrological alteration; contains few roads or buildings; noxious weed or ANVS cover is ≤30%.						
AA cultivated or heavily grazed or logged; subject to relatively substantial fill placement, grading, clearing, or hydrological alteration; high road or building density; or noxious weed or ANVS cover is >30%.						

Comments	(types c	of disturbance,	intensity,	season	etc.):	
----------	----------	-----------------	------------	--------	--------	--

13. STRUCTURAL DIVERSITY (Based on number of "Cowardin" vegetated classes present [do not include unvegetated classes]; see #10 above.)

Existing # of "Cowardin" Vegetated Classes in AA	Initial Rating	Is current management peristence of additional		Modified Rating
≥3 (or 2 if one is forested) classes		NA	NA	NA
2 (or 1 if forested) classes	mod	NA	NA	NA
1 class, but not a monoculture		←NO	YES→	
1 class, monoculture (1 species comprises ≥90% of total cover)		NA	NA	NA

Comments:	
Comments:	

ii. Prominent noxious, aquatic nuisance, and other exotic vegetation species: ____

iii. Provide brief descriptive summary of AA and surrounding land use/habitat: AA is large wet meadow, emergent marsh and shrub/scrub wetland that has been restored by plugging man-made drain ditches. AA is not grazed or hayed. Surrounding land is moderately grazed and managed for agriculture.

14A. HABITAT FOR FEDER	ALLY	LISTE	D OR	PRO	POSE	D THE	REATE	NED	OR E	NDAN	GERE	D PL	ANTS	OR A	NIMAL	.s				
Primary or critical habitat (I Secondary habitat (Iist spe	AA is Documented (D) or Suspected (S) to contain: Check box based on definitions in manual. Primary or critical habitat (list species)																			
ii. Rating: Based on the strong	ngest h	abitat	chose	n in '	14A(i) a	above,	selec	t the	corres	pondin	g func	tiona	point	and ra	ting.					_
Highest Habitat Level	Doc/F	Primar	y S	us/P	rimary	Do	c/Sec	onda	ry S	us/Se	conda	ry	Doc/In	ciden	tal	Sus/Incidental None				
Functional Point/Rating				-						-									0L	
Sources for documented us	e (e.g.	obser	vation	s, red	cords):		_													
	14B. HABITAT FOR PLANTS OR ANIMALS RATED S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM Do not include species listed in 14A above.																			
i. AA is Documented (D) or Suspected (S) to contain: Check box based on definitions in manual. Primary or critical habitat (list species)																				
ii. Rating: Based on the stro	ngest l	habita	t chose	en in	14A(i)	above	, selec	t the	corres	pondii	ng fund	ctiona	ıl point	and ra	ating.					_
Highest Habitat Level	Doc/F	rimar	y S	us/P	rimary	Do	c/Sec	onda	ry S	us/Se	conda	ry	Doc/In	ciden	tal S	Sus/I	ncider	ntal	None	
S1 Species Functional Point/Rating				-						-										
S2 and S3 Species Functional Point/Rating				_						-				.2L			.1L			
Sources for documented us	e (e n	ohsei	vation	s red	ords).	Grav	wolf do	CUM	ented	onsite	hy land	down	er acco	ordina	to MD	Т				J
i. Evidence of Overall Wildl Substantial: Based on an observations of abunda abundant wildlife sign of presence of extremely interview with local biology. Moderate: Based on any of observations of scatters common occurrence of	14C. GENERAL WILDLIFE HABITAT RATING i. Evidence of Overall Wildlife Use in the AA: Check substantial, moderate, or low based on supporting evidence. Substantial: Based on any of the following [check]. Minimal: Based on any of the following [check]. few or no wildlife observations during peak use periods little to no wildlife sign such as scat, tracks, nest structures, game trails, etc. little to no wildlife sign sparse adjacent upland food sources interview with local biologist with knowledge of the AA Moderate: Based on any of the following [check]. Subservations of scattered wildlife groups or individuals or relatively few species during peak periods common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. adequate adjacent upland food sources common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc. commo																			
ii. Wildlife Habitat Features For class cover to be conside percent composition of the AA S/I = seasonal/intermittent; T/	red eve \ (see #	enly di #10).	stribute Abbrev	ed, th viatio	ne mos ns for s	t and I surface	east p	reval r dura	ent ve g ations	getate are as	d class follows	ses m s: P/F	nust be P = per	within maner	20% ont/pere	of ead nnial	ch othe			
Structural Diversity					High								derate						ow	
(see #13) Class Cover Distribution (all vegetated classes)		E	ven			☐ Un	even			E				⊠ Un	even			E		
Duration of Surface Water in ≥ 10% of AA	P/P	S/I	T/E	Α	P/P	S/I	T/E	Α	P/P	S/I	T/E	Α	P/P	S/I	T/E	Α	P/P	S/I	T/E	Α
														Н						
(see #12i)														"						
☐ Moderate Disturbance at AA (see #12i)																				
☐ High Disturbance at AA (see #12i)																				
iii. Rating: Use the conclusi	ons fro	m i an	d ii ab	ove a	and the	matri	x belov	v to s	elect t	he fun	ctional	poin	t and ra	ating.						
Evidence of Wildlife Use									ature											
(i)		Exc	eptio	nal		\boxtimes	High			☐ Mc	derate)		☐ Lo	W					
Substantial	_																			
							7M			-						-				
Minimal Commentary various bird and	nino ==	o.m.:	 lo bar		0.00.0	*00	th sign	.:£:	A 14-11-11	lifa be	hitot					_				
Comments: various bird spec	cies, m	amma	ıs, her	ps. L	_arge a	rea w	ith sigr	nificai	nt Wildl	iite ha	oitat.									

		Wetland/Site #(s): al wetlands outside Rock Creek corridor																
14D. GENERAL FISH HABIT If the AA is not used by entrapped in a canal], the	fish, fisl	n use is n		able du	e to h		const	raints,	or is n	ot desi	ired fro	om a r	nanag	ement	perspe	ective	(such a	as fish
Assess this function if the precluded by perched co		,		e existi	ng sit	tuation i	s "co	rrecta	ble" su	ch that	t the A	A cou	ld be ι	used by	/ fish [i	.e., fis	h use	is
Type of Fishery: Co	old Wat	er (CW)	☐ War	m Wate	er (W	W) Us	e the	CW o	r WW	guideli	nes in	the ma	anual t	o comp	lete th	e matr	ix.	
. Habitat Quality and Know	n / Sus _l	pected F	ish Spec	ies in <i>l</i>	AA:	Use ma	trix to	sele	ct the fu	unction	nal poi	nt and	rating	J.				_
Duration of Surface Water in AA	☐ Pe	rmanent	/ Perenr	nial		☐ Se	asor	al / Ir	ntermit	tent		□т	empo	rary / E	Ephen	neral		
Aquatic Hiding / Resting /]													
Escape Cover Thermal Cover:	Opti	mal Ac	dequate	Po	or	Optir	nal	Ade	quate	Po	or	Opt	imal	Adec	uate	Po	or	
optimal / suboptimal	0	S	S	0	S	0	S	0	S	0	S	0	S	0	S	0	S	
FWP Tier I fish species																		
FWP Tier II or Native																		
Game fish species FWP Tier III or Introduced																		
Game fish																		
FWP Non-Game Tier IV or No fish species																		
Sources used for identifying	fish sp	p. poter	tially for	und in	AA:													
ii. Modified Rating: NOTE: N	Modified	d score ca	annot exc	eed 1.0	or b	e less t	han ().1.										
a) Is fish use of the AA signific MDEQ list of waterbodies in ne																		
support, or do aquatic nuisand																		
b) Does the AA contain a docu native fish or introduced game										nctuary	pool,	upwe	lling ai	rea; sp	ecify in	ı comr	nents)	for
ii. Final Score and Rating:	Com	ments: _																
14E. FLOOD ATTENUATION Applies only to wetlands If wetlands in AA are no	that ar	e subject	(procee to floodi channel	ng via ii	n-cha	innel or flow, cl	over neck	bank t	flow. A box a	ınd pro	oceed	to 14F	·.					
Entrenchment Ratio (ER) Est Flood-prone width = estimated																	e of the	e strean
/	=						0	L.							A	TOPE .		
flood prone width / bankfull wid	dth = en	trenchme	ent ratio		2	D 16		OH THE STATE OF TH	Milial	Ye way	kd.	- _		Jan.	– Æ	lood-p	rone W	'idth
					23	k Bankfu	III De	pth 🦠		i	***********	<u>.</u>	<u>"</u>		Bank	cfull W	idth	
									В	ankfull	Depth	Vouvoc	wolf .					
Slightly Entr	enched	1		Mode	erate	ly Entre	ench	ed				Fnti	renche	ed he				
ER ≥ 2	.2				ER =	1.41 – 2	2.2				1	ER =	1.0 –	1.4	_			
C stream type D stream t		E strear	n type		B str	eam typ	e		A stre	am typ	oe 1	Fstr	eam ty	ype	G str	ream t	уре	
					1		1					Ę		-5		/		
										·								
i. Rating: Working from top to	bottom	n. use the	matrix b	elow to	seled	ct the fu	nctio	nal po	oint and	rating	ı.							
Estimated or Calculated (Rosgen 1994, 1996)			□S	lightly E	ntrer	nched		Mode	erately stream	Entren				Entrend Strear		e e		
Percent of Flooded Wetland		ified as]													
Forested and/or Scrub/Sh AA contains no outlet or re		l outlot	75%	25-7		<25%	_	5%	25-75	% <	<25%	75		25-759	% <	25%		
AA contains no outlet or re							-											
AA GOIRAINS UNI CSUIC	ou ouli						- 11										Ī	

ii. Are ≥10 acres of wetland in the AA subject to flooding AND are man-made features which may be significantly damaged by floods located
within 0.5 mile downstream of the AA? TYES NO Comments:

14F.	SHORT AND LONG TERM SURFACE WATER STORAGE	☐ NA (proceed to 14G)
	Applies to wetlands that flood or pond from overbank or in-chann	nel flow, precipitation, upland surface flow, or groundwater flow.
	If no wetlands in the AA are subject to flooding or ponding, then	check the NA box and proceed to 14G.

i. Rating: Working from top to bottom, use the matrix below to select the functional point and rating. Abbreviations for surface water durations are as follows: P/P = permanent/perennial; S/I = seasonal/intermittent; and T/E = temporary/ephemeral [see manual for further definitions of these terms].

Estimated Maximum Acre Feet of Water Contained in Wetlands within the AA that are Subject to Periodic Flooding or Ponding		>5 acre fe	eet	□ 1.1	to 5 ac	re feet	☐ ≤1 acre foot		
Duration of Surface Water at Wetlands within the AA	⊠ P/P	□ S/I	□ T/E	□ P/P	□ S/I	□ T/E	□ P/P	□ S/I	□ T/E
Wetlands in AA flood or pond ≥ 5 out of 10 years	1H								
Wetlands in AA flood or pond < 5 out of 10 years									

^			4	۱.	_
u	mc	me	m	IS	:

14G. SEDIMENT / NUTRIENT / TOXICANT / RETENTION AND REMOV	AL NA (proceed to 14H
---	-----------------------

Applies to wetland with potential to receive sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, check the NA box and proceed to 14H.

i. Rating: Working from top to bottom, use the matrix below to select the functional point and rating.

Sediment, Nutrient, and Toxicant Input Levels within AA	AA receive has potent nutrients, such that a substantia sedimenta toxicants, present.	tial to deliv or compou other funct illy impaire tion, sourc	er sedime nds at lev ions are n d. Minor es of nutr	ents, rels not rients or	Waterbody is on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.						
% Cover of Wetland Vegetation in AA	⊠≥′	70%	□ < 70%		□ ≥ 70 %		□ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70% □ < 70%				
Evidence of Flooding / Ponding in AA	⊠ Yes	☐ No	☐ Yes	☐ No	☐ Yes	□No	☐ Yes	☐ No			
AA contains no or restricted outlet	1H										
AA contains unrestricted outlet											

	ents	

14H. SEDIMENT / SHORELINE STABILIZATION NA (proceed to 14l)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body which is subject to wave action.

If 14H does not apply, check the NA box and proceed to 14I.

% Cover of Wetland Streambank or Shoreline by Species with Stability	Duration of S	Duration of Surface Water Adjacent to Rooted Vegetation						
Ratings of ≥6 (see Appendix F).	☐ Permanent / Perennial	☐ Temporary / Ephemera						
□ ≥ 65%								
□ 35-64%								

Comments:

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. Level of Biological Activity: Synthesis of wildlife and fish habitat rates (select).

General Fish Habitat Rating	General Wildlife Habitat Rating (14Ciii)							
(14Diii)	□ E/H	⊠ M	□ L					
☐ E/H								
								
⊠ NA		M						

ii. Rating: Working from top to bottom, use the matrix below to select the functional point and rating. Factor A = acreage of vegetated wetland component in the AA; Factor B = level of biological activity rating from above (14li); Factor C = whether or not the AA contains a surface or subsurface outlet; the final three rows pertain to the duration of surface water in the AA, where P/P, S/I, and T/E were previously defined, and A = "absent" [see manual for further definitions of these terms].

Α	☑ Vegetated Component >5 acres				☐ Vegetated Component 1-5 acres					☐ Vegetated Component <1 acre								
В		ligh	⊠M	oderate		Low	_ _	ligh	☐ Mo	derate		Low	_ 	ligh	☐ Mo	derate	L	.ow
С	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No	Yes	No
P/P				.5M														
S/I																		
T/E/A																		

					s): <u>ali wetiand</u>				
14I. PRODUCTION EXPORT / FOOD (CHAIN S	UPPORT (con	itinued)						
iii. Modified Rating: Note: Modified so	ore cann	not exceed 1.0	or be less that	n 0.1.					
Vegetated Upland Buffer: Area with ≥ 30% plant cover, ≤ 15% noxious weed or ANVS cover, AND that is not subjected to periodic me mowing or clearing (unless for weed control). Is there an average ≥ 50-foot wide vegetated upland buffer around ≥ 75% of the AA's perimeter? YES, add 0.1 to score in ii = 0.6									
iv. Final Score and Rating: $\underline{.6M}$ Con	nments:								
14J. GROUNDWATER DISCHARGE / Check the appropriate indicators i									
i. Discharge Indicators The AA is a slope wetland. Springs or seeps are known Vegetation growing during of wetland occurs at the toe of Seeps are present at the word AA permanently flooded du Wetland contains an outlet, Shallow water table and the Other:	dormant of a natur etland ed ring drou but no in	season/drougl ral slope. dge. ught periods. nlet.	ht.	☐ Pe ☐ We ☐ Sti	etland contain	ors strate present v is inlet but no d wn 'losing' stre	outlet.	, , ,	0 ,
iii. Rating: Use the information from i a	and ii abo	ove and the tal	ole below to se	lect the fu	unctional poin	t and rating.			
Criteria		Duration of Saturation at AA Wetlands FROM GROUNDWATER DISCHARGE or WITH WATER THAT IS RECHARGING THE GROUNDWATER SYSTEM ☑ P/P ☐ S/I ☐ T ☐ None							
☐ Groundwater Discharge or Recha	arge	1H							
☐ Insufficient Data/Information	g -		L	1					
Comments:									
i. Rating: Working from top to bottom,	AA co	ntains fen, bo	g, warm	AA doe	s not contai	n previously	AA doe	es not contai	n
	AA co	ntains fen, bo is or mature (: ed wetland Of iation listed a	og, warm >80 yr-old) ⋜ plant	AA doe cited ra diversi contair		n previously O structural gh OR ciation	previou associ	es not contai usly cited rar ations AND s ty (#13) is lo	e types OR structural
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11)	AA conspring forester associ	ntains fen, bo s or mature (; ed wetland Of iation listed a INHP	og, warm >80 yr-old) ⋜ plant	AA doe cited ra diversi contair listed a	es not contain are types ANI ty (#13) is hig as plant asso	n previously O structural gh OR ciation e MTNHP	previou associ diversi	usly cited rar ations AND s	re types OR structural w-moderate
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11) Low Disturbance at AA (#12i)	AA conspring forester associathe MT	ntains fen, bo s or mature (; ed wetland Of iation listed a INHP	og, warm >80 yr-old) ⋜ plant s "S1" by	AA doe cited ra diversi contair listed a	es not contain are types ANI ty (#13) is high as plant asso as "S2" by the	n previously O structural gh OR ciation e MTNHP	previou associ diversi	usly cited rar ations AND s ty (#13) is lo	re types OR structural w-moderate
Replacement Potential Estimated Relative Abundance (#11) Low Disturbance at AA (#12i) Moderate Disturbance at AA (#12i)	AA colspring foreste associ the M1	ntains fen, bo is or mature (: ed wetland Of iation listed a TNHP © Common .9H	og, warm >80 yr-old) R plant s "S1" by	AA doe cited ra diversi contair listed a	es not contain are types ANI ty (#13) is high as plant asso as "S2" by the Common	n previously D structural gh OR ciation e MTNHP Abundant	previou associ diversi Rare	usly cited rar ations AND s ty (#13) is lo	e types OR structural w-moderate Abundant
Replacement Potential Estimated Relative Abundance (#11) Low Disturbance at AA (#12i) Moderate Disturbance at AA (#12i) High Disturbance at AA (#12i)	AA conspring forester associate MT	ntains fen, bo is or mature (: ed wetland Of iation listed a FNHP	og, warm >80 yr-old) R plant s "S1" by	AA doe cited ra diversi contair listed a	es not contain are types ANI ty (#13) is high as plant asso as "S2" by the Common	n previously D structural gh OR ciation e MTNHP Abundant	previou associ diversi	usly cited rar ations AND s ty (#13) is lo	e types OR structural w-moderate Abundant
Replacement Potential Estimated Relative Abundance (#11) Low Disturbance at AA (#12i) Moderate Disturbance at AA (#12i)	AA cospring foreste associthe M1 Rare	ntains fen, bo s or mature (: ed wetland Of iation listed a FNHP SOCOMMON 9H eational or edu or education	og, warm >80 yr-old) R plant s "S1" by Abundant NA (proceed acational oppor	AA doe cited ra diversi contair listed a Rare to Overa tunity.	es not container types ANI ty (#13) is high as plant asso as "S2" by the Common Il Summary ani ii. NO, cl	n previously D structural gh OR ciation e MTNHP D Abundant nd Rating page	previous associ diversi Rare	usly cited rar ations AND s ty (#13) is lo	e types OR structural w-moderate Abundant
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11) Low Disturbance at AA (#12i) Moderate Disturbance at AA (#12i) High Disturbance at AA (#12i) Comments: Fen present. 14L. RECREATION / EDUCATION POAffords 'bonus' points if AA provide i. Is the AA a known or potential recreii. Check categories that apply to the	AA co spring foreste associ the M1 Rare	ntains fen, bo s or mature (: ed wetland Of iation listed a TNHP Common 9H eational or edu or education Educational/S Other:	og, warm >80 yr-old) R plant s "S1" by Abundant NA (proceed acational opporal site? YE	AA doe cited ra diversi contair listed a Rare to Overa tunity.	es not container types ANI ty (#13) is high as plant asso as "S2" by the Common Il Summary ani ii. NO, cl	n previously D structural gh OR ciation e MTNHP D Abundant nd Rating page	previous associ diversi Rare	usly cited rar ations AND s ty (#13) is lo	e types OR structural w-moderate Abundant
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11) □ Low Disturbance at AA (#12i) □ Moderate Disturbance at AA (#12i) □ High Disturbance at AA (#12i) Comments: Fen present. 14L. RECREATION / EDUCATION PO Affords 'bonus' points if AA provide i. Is the AA a known or potential recreii. Check categories that apply to the iii. Rating: Use the matrix below to select	AA coi spring foreste associ the M1 Rare	ntains fen, bo s or mature (sed wetland Officiation listed a TNHP Solution Common 9H eational or education Educational/Solutional point	og, warm >80 yr-old) R plant s "S1" by Abundant NA (proceed acational opporal site? YE) Scientific Study and rating.	AA doe cited ra diversi contain listed a Rare	es not container types ANI ty (#13) is high as plant asso as "S2" by the Common Il Summary ani ii. NO, cl	n previously D structural gh OR ciation e MTNHP D Abundant nd Rating page	previous associativersi	usly cited rar ations AND s ty (#13) is lor Common	e types OR structural w-moderate Abundant
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11) Low Disturbance at AA (#12i) Moderate Disturbance at AA (#12i) High Disturbance at AA (#12i) Comments: Fen present. 14L. RECREATION / EDUCATION POAffords 'bonus' points if AA provide i. Is the AA a known or potential recreii. Check categories that apply to the liii. Rating: Use the matrix below to select the selection of the s	AA column spring forester associate MT Rare	ntains fen, bo s or mature (sed wetland Officiation listed a TNHP Solution Common 9H eational or education Educational/Solutional point Recreational	og, warm >80 yr-old) R plant s "S1" by Abundant NA (proceed acational opporal site? YES Scientific Study and rating.	AA doe cited ra diversi contain listed a Rare to Overatunity. S, go to	es not container types ANI ty (#13) is high as plant asso as "S2" by the Common Ill Summary ani ii. NO, classumptive Recomments	n previously D structural gh OR ciation e MTNHP D Abundant nd Rating page	previous associ diversi Rare	usly cited rar ations AND s ty (#13) is lor Common	e types OR structural w-moderate Abundant
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11) □ Low Disturbance at AA (#12i) □ Moderate Disturbance at AA (#12i) □ High Disturbance at AA (#12i) Comments: Fen present. 14L. RECREATION / EDUCATION PO Affords 'bonus' points if AA provide i. Is the AA a known or potential recreii. Check categories that apply to the iii. Rating: Use the matrix below to select	AA cospring forester associate MT Rare	ntains fen, bo s or mature (sed wetland Officiation listed a TNHP Solution Common 9H eational or education Educational/Solutional point Recreational eneral public	og, warm >80 yr-old) R plant s "S1" by Abundant NA (proceed acational opporal site? YE Scientific Study and rating. I or Education access (no p	AA doe cited ra diversi contain listed a Rare to Overatunity. S, go to Cor	es not container types ANI ty (#13) is high as plant asso as "S2" by the Common Ill Summary ani ii. NO, classumptive Recomments	n previously D structural gh OR ciation e MTNHP D Abundant nd Rating page	previous associativersi Rare	usly cited rar ations AND s ty (#13) is lor Common	e types OR structural w-moderate Abundant
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11) □ Low Disturbance at AA (#12i) □ Moderate Disturbance at AA (#12i) □ High Disturbance at AA (#12i) Comments: Fen present. 14L. RECREATION / EDUCATION PO Affords 'bonus' points if AA provide i. Is the AA a known or potential recreii. Check categories that apply to the iii. Rating: Use the matrix below to select Known or Public ownership or public easemer	AA cospring foreste associthe M1 Rare	ntains fen, bo s or mature (: ed wetland Of iation listed a rNHP Sommon 9H eational or edu or education Educational/S Other: nctional point Recreational eneral public s (no permis	og, warm >80 yr-old) R plant s "S1" by Abundant NA (proceed acational oppor al site? YE Scientific Study and rating. I or Education access (no p	to Overatunity. S, go to Cal Area ermissio	ss not container types ANI ty (#13) is high s plant asso s "S2" by the Common Ill Summary ani ii. NO, clusumptive Reconstruction	n previously D structural gh OR ciation e MTNHP D Abundant and Rating page	previous associdiversi la Rare	usly cited rar ations AND s ty (#13) is lor Common sumptive recr	e types OR structural w-moderate Abundant
i. Rating: Working from top to bottom, Replacement Potential Estimated Relative Abundance (#11) □ Low Disturbance at AA (#12i) □ Moderate Disturbance at AA (#12i) □ High Disturbance at AA (#12i) Comments: Fen present. 14L. RECREATION / EDUCATION PO Affords 'bonus' points if AA provide i. Is the AA a known or potential recreii. Check categories that apply to the iii. Rating: Use the matrix below to select Known or Public ownership or public easemer Private ownership with general publice.	AA cospring foreste associthe M1 Rare	ntains fen, bo s or mature (: ed wetland Of iation listed a rNHP Sommon 9H eational or edu or education Educational/S Other: nctional point Recreational eneral public s (no permis	og, warm >80 yr-old) R plant s "S1" by Abundant NA (proceed acational oppor al site? YE Scientific Study and rating. I or Education access (no p	to Overatunity. S, go to Cal Area ermissio	ss not container types ANI ty (#13) is high s plant asso s "S2" by the Common Ill Summary ani ii. NO, clusumptive Reconstruction	n previously D structural gh OR ciation e MTNHP D Abundant and Rating page	previous associativersi	suly cited rar ations AND s ty (#13) is lor Common sumptive recr	e types OR structural w-moderate Abundant

Function & Value Variables	Rating – Actual Functional Points	Possible Functional Points	Functional Units: Actual Points x Estimated AA Acreage	Indicate the Four Most Prominent Functions with an Asterisk
A. Listed / Proposed T&E Species Habitat	low 0.00	1.00	0	
B. MT Natural Heritage Program Species Habitat	low 0.20	1.00	7.96	
C. General Wildlife Habitat	mod 0.70	1.00	27.87	
D. General Fish Habitat	NA	NA		
E. Flood Attenuation	NA	NA		
F. Short and Long Term Surface Water Storage	high 1.00	1.00	39.81	
G. Sediment / Nutrient / Toxicant Removal	high 1.00	1.00	39.81	
H. Sediment / Shoreline Stabilization	NA	NA		
I. Production Export / Food Chain Support	mod 0.60	1.00	23.89	
J. Groundwater Discharge / Recharge	high 1.00	1.00	39.81	
K. Uniqueness	high 0.90	1.00	35.83	
L. Recreation / Education Potential (bonus point)	low 0.05		1.99	
Total Points	5.45	8	217 Total	Functional Units
Percent of Possibl	e Score 68% (round	to nearest whol	e number)	

Category I Wetland: (must satisfy one of the following criteria; otherwise go to Category II) Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; or Score of 1 functional point for Uniqueness; or Score of 1 functional point for Flood Attenuation and answer to Question 14E.ii is "yes"; or Percent of possible score > 80% (round to nearest whole #).
Category II Wetland: (Criteria for Category I not satisfied and meets any one of the following criteria; otherwise go to Category IV) Score of 1 functional point for MT Natural Heritage Program Species Habitat; or Score of .9 or 1 functional point for General Wildlife Habitat; or Score of .9 or 1 functional point for General Fish Habitat; or "High" to "Exceptional" ratings for both General Wildlife Habitat and General Fish/Aquatic Habitat; or Score of .9 functional point for Uniqueness; or Percent of possible score > 65% (round to nearest whole #).
☐ Category III Wetland: (Criteria for Categories I, II, or IV not satisfied)
Category IV Wetland: (Criteria for Categories I or II are not satisfied and all of the following criteria are met; if not go to Category III) "Low" rating for Uniqueness; and Vegetated wetland component < 1 acre (do not include upland vegetated buffer); and Percent of possible score < 35% (round to nearest whole #).
OVERALL ANALYSIS AREA (AA) RATING: Check the appropriate category based on the criteria outlined above.

Appendix C

2009 REPRESENTATIVE PHOTOGRAPHS

MDT Wetland Mitigation Monitoring Big Hole Grazing Association Beaverhead County, Montana



Photo 1: PP1 – View facing north.



Photo 2: PP1 – View facing west.



Photo 3: PP1 – View facing east.



Photo 4: PP2 – View facing northwest.



Photo 5: PP2 – View facing west.



Photo 6: PP2 – View facing east.



Photo 7: PP3 – View facing southeast.



Photo 8: PP3 – View facing west.



Photo 9: PP3 – View facing northwest.



Photo 10: PP4 – View facing southeast.



Photo 11: PP4 – View facing southeast.



Photo 12: PP4 – View facing southwest.



Photo 13: PP5 - View facing east.



Photo 14: PP5 - View facing southwest.



Photo 15: PP5 - View facing northwest.



Photo 16: PP6 - View facing southeast.



Photo 17: PP6 - View facing south.



Photo 18: PP6 - View facing northwest.



Photo 19: PP7 - View facing southwest.



Photo 20: PP7 - View facing west.



Photo 21: PP7 - View facing south.



Photo 22: Transect Start.



Photo 23: Transect End.




Photo 24: Moose in the site.

Appendix D

PROJECT PLAN SHEET

MDT Wetland Mitigation Monitoring Big Hole Grazing Association Beaverhead County, Montana



Appendix E

BIRD SURVEY PROTOCOL GPS PROTOCOL

MDT Wetland Mitigation Monitoring Big Hole Grazing Association Beaverhead County, Montana

BIRD SURVEY PROTOCOL

This protocol was developed by the Montana Department of Transportation (MDT) to monitor bird use within their Wetland Mitigation Sites. Though each wetland mitigation site is vastly different, the bird survey data collection methods were standardized to order to increase repeatability. The protocol uses an "area search within a restricted time frame" to collect data on bird species, density, behavior, and habitat-type use.

Survey Area

Sites that can be entirely walked: Sites where the entire perimeter or area can be walked include, but are not limited to: small ponds, enhanced historic river channels, and wet meadows. If the wetland is not uncomfortably inundated, walk several meandering transects to sufficiently cover the wetland. Meandering transects can be used, even if a small portion of the area is inaccessible (e.g. cannot cross due to inundation). Use binoculars to identify the bird species, to count the number of individuals, and to identify their behavior and habitat type. Data can be recorded directly onto the bird survey form or into a field notebook. The number of meandering transects and their direction (or location) should be recorded in the field notebook and/or drawn onto the aerial photograph or topographic map. Meandering transects are not formal and should not be staked. Each site should be walked and surveyed to the fullest extent within the set time limit.

Sites than cannot be entirely walked: Sites where the entire perimeter or area cannot be walked include, but are not limited to: very large sites (i.e. perimeter of 2-3 miles), and large-bodied waters (i.e. reservoirs), where deep water habitat (> 6 feet) is close to shore. For large-bodied waters where only one area was graded to create or enhance the development of wetland, bird surveys should be walked along meandering transects within or around the graded area (see above.). For sites that cannot be walked, bird surveys should be conducted from many lookout posts, established at key vantage points. The general location of lookout posts should be recorded in the field notebook or drawn onto the aerial photograph or topographic map. Lookout post locations do not need to be staked. Both binoculars and spotting scopes may be used in order to accurately identify and count the birds. Depending upon the size of the open water, more time may be spent viewing the mitigation area from lookout posts than is spent traveling between posts.

Survey Time

Ideally, bird surveys should be conducted in the morning hours when bird activity is often greatest (i.e. sunrise to no later than 11:00 am). Surveys can be completed before 11am if all transects have been walked or all lookout posts have been viewed with no new bird activity observed. For some sites bird surveys may need to be performed in the late afternoon or evening due to traveling constraints or weather. The overall limiting time factor will be the number of budgeted hours for the project.

Data Recording

Bird Species List: Record each bird species observed onto the Bird Survey-Field Data Sheet (or field notebook). Record the bird's common name using the appropriate 4-letter code. The 4-letter code uses the first two letters of the first two word's of the bird's common name or if one name, the first four letters. For example, Mourning Dove is coded as MODO while Mallard is coded as MALL. If an unknown individual is observed, use the 4-letter protocol, but define your

PBS

BIRD SURVEY PROTOCOL (continued)

abbreviation at the bottom of the field data sheet. For example, unknown shorebird is UNSB; unknown brown bird is UNBR; unknown warbler is UNWA; and unknown waterfowl is UNWF. For a flyover of a flock of unknown species, use a term that describes the birds' general characteristics and include the approximate flock size in parenthesis; do not fill in the habitat column. For example, a flock of black, medium-sized birds could be coded as UNBB / FO (25).

Bird Density: For each observation record the actual or estimated number of individuals observed per species and per behavior. Totals can be tallied in the office and entered onto the Bird Survey-Field Data Sheet.

Bird Behavior: Bird behavior must be identified by what is known. When a species is observed, the behavior that is immediately exhibited is recorded. Only behaviors that have discreet descriptive terms should be used. The following terms are recommended: breeding pair (BP); foraging (F); flyover (FO); loafing (L), which is defined as sleeping, roosting, or floating with head tucked under wing; and nesting (N). If other behaviors that have a specific descriptive word are observed then it can be used and should later be added to the protocol. Descriptive words or phrases such as "migrating" or "living on site" are unknown behaviors.

Bird Species Habitat Use: When a species is observed, the habitat is also recorded. The following broad habitat categories are used:

- aquatic bed (AB), defined as rooted-floating, floating-leaved, or submergent vegetation.
- marsh (MA), defined as emergent (e.g. cattail, bulrush) vegetation with surface water.
- wet meadow (WM), defined as grasses, sedges, or rushes with little to no surface water.
- scrub-shrub (SS), defined as shrub covered wetland.
- forested (FO), defined as tree covered wetland.
- open water (OW), defined as unvegetated surface water.
- upland (UP), defined as the upland buffer.

Other categories can be used and defined on the data sheet and should later be added to the protocol.

Other Fields

Bird Visit: Each bird survey (i.e. spring, fall, and mid-season) should be completed on separate Bird Survey-Field Data Sheets.

Time: Record the start time and end time on the Bird Survey-Field Data Sheet.

Date: Record the date of the bird survey.

Weather: Record the weather conditions (i.e. temperature, wind, condition).

Notes: Note if a particular individual bird is using a constructed nest box and note the condition of constructed nest box(es). Also record any comments about the site, wildlife, wetland conditions, etc.



GPS MAPPING AND AERIAL PHOTO REFERENCING PROCEDURE

From 2001 through 2006, PBS&J mapped the vegetation community boundaries, photograph points, and other sampling locations in the field using the resource-grade Trimble GEO III GPS (Global Positioning System) unit. The data were collected with a minimum of three positions per feature using Course/Acquisition code. The collected data were then transferred to a personal computer (PC) and differentially corrected to the nearest operating Community Base Station. The corrected data were then exported to ACAD drawings in Montana State Plain Coordinates NAD 83 international feet. The Trimble GEO III GPS unit was also used for some sites in 2007.

The collected and processed Trimble Geo III GPS positions had a 68% accuracy of 7 feet except in isolated areas where accuracy fell to 12 feet. This is within the 1 to 5 meter range listed as the expected accuracy of the mapping grade Trimble GPS.

In 2007 and 2008 sites were mapped using the resource-grade Magellan MobileMapper Office GPS unit. The Magellan GPS unit has a comparable accuracy level to the Trimble Geo III unit.

Each year, MDT photographs each mitigation site from the air. These aerial photographs are not geo-referenced, but serve as a visual aid to map wetland development and vegetation communities, and to show approximate locations for various monitoring activities (i.e. photograph points, transects, or macroinvertebrate sampling). Reference points that are observable on the aerial photo (i.e. road, stream channel, or fence) were also marked with the GPS unit in order to better position the aerial photograph. This positioning did not remove any of the distortion inherent to all photos. All mapped features and community boundaries were reviewed by the wetland biologist, to increase the figure's accuracy.

Any relationship of features located to easement or property lines are not to be construed from these figures. These relationships can only be determined with a survey by a licensed surveyor.

