
MONTANA DEPARTMENT OF TRANSPORTATION WETLAND MITIGATION MONITORING REPORT: YEAR 2004

*Big Spring Creek
Lewistown, Montana*



Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Avenue
Helena, MT 59620-1001

Prepared by:

LAND & WATER CONSULTING
~ A DIVISION OF **PBS&J**
P.O. Box 239
Helena, MT 59624

June 2005

Project No: B43054.00 - 0502



MONTANA DEPARTMENT OF TRANSPORTATION

WETLAND MITIGATION MONITORING REPORT:

YEAR 2004

*Big Spring Creek
Lewistown, Montana*

Prepared for:

MONTANA DEPARTMENT OF TRANSPORTATION
2701 Prospect Ave
Helena, MT 59620-1001

Prepared by:

LAND & WATER CONSULTING
A DIVISION OF PBS&J
P.O. Box 239
Helena, MT 59624

June 2005

Project No: B43054.00 - 0502



TABLE OF CONTENTS

1.0 INTRODUCTION.....	1
2.0 METHODS.....	3
2.1 Monitoring Dates and Activities.....	3
2.2 Hydrology.....	3
2.3 Vegetation.....	4
2.4 Soils.....	4
2.5 Wetland Delineation.....	5
2.6 Mammals, Reptiles, and Amphibians.....	5
2.7 Birds.....	5
2.8 Macroinvertebrates.....	5
2.9 Functional Assessment.....	6
2.10 Photographs.....	6
2.11 GPS Data.....	6
2.12 Maintenance Needs.....	6
3.0 RESULTS.....	6
3.1 Hydrology.....	6
3.2 Vegetation.....	7
3.3 Soils.....	12
3.4 Wetland Delineation.....	12
3.5 Wildlife and Fish.....	12
3.6 Macroinvertebrates.....	13
3.7 Functional Assessment.....	14
3.8 Photographs.....	15
3.9 Maintenance Needs/Recommendations.....	15
3.10 Current Credit Summary.....	17
4.0 REFERENCES.....	18

TABLES

Table 1	<i>2001 - 2004 Big Spring Creek vegetation species list.</i>
Table 2	<i>Vegetation transect data summary.</i>
Table 3	<i>2004 observed mortality of planted woody species.</i>
Table 4	<i>Fish and wildlife species observed on the Big Spring Creek Mitigation Site 2001-2004</i>
Table 5	<i>Summary of 2004 wetland function/value ratings and functional points at the Big Spring Creek Mitigation Project.</i>

FIGURES

Figure 1	<i>Project Site Location Map</i>
Figure 2	<i>Monitoring Activity Locations 2004</i>
Figure 3	<i>Mapped Site Features 2004</i>

CHARTS

Chart 1	<i>Transect maps showing vegetation types from the start of transect (0 feet) to the end of transect (418 feet) for each year monitored.</i>
Chart 2	<i>Length of vegetation communities along Transect 1.</i>
Chart 3	<i>Macroinvertebrate bioassessment scores 2001 – 2004.</i>

APPENDICES

Appendix A	<i>Figures 2 & 3</i>
Appendix B	<i>Completed 2004 Wetland Mitigation Site Monitoring Form</i> <i>Completed 2004 Bird Survey Forms</i> <i>Completed 2004 Wetland Delineation Forms</i> <i>Completed 2004 Functional Assessment Forms</i>
Appendix C	<i>Representative Photographs, 2004 Aerial Photograph</i>
Appendix D	<i>Conceptual Site Layout</i>
Appendix E	<i>Bird Survey Protocol</i> <i>GPS Protocol</i>
Appendix F	<i>Macroinvertebrate Sampling Protocol and Data</i>

1.0 INTRODUCTION

In 1996, the Montana Fish, Wildlife & Parks (FWP) approached the Montana Department of Transportation (MDT) with a partnership proposal to restore approximately 0.5 mile of Big Spring Creek, at the FWP Brewery Flats Fishing Access site, 1 mile SE of Lewistown in Fergus County (**Figure 1**). Big Spring Creek was straightened through the Brewery Flats area around 1907 by the Milwaukee Railroad to facilitate the construction of a freight yard to the west of the creek. FWP proposed, through their Future Fisheries Improvement Program (FFIP), to restore that section of Big Spring Creek that traversed Brewery Flats to a more natural condition for the purpose of improving fisheries habitat. In addition to increasing total stream length from 2,300 feet to 4,000 feet, the design also included the establishment of a functional floodplain and associated wetland habitat.






In 1998, an MOA between MDT and FWP was signed by the agencies, thus formalizing a cooperative agreement to restore Big Spring Creek. In return for a cash contribution to the project, MDT would receive 7.21 acres of Corps of Engineers (COE)-approved wetland mitigation credit to provide mitigation for projected wetland impacts resulting from MDT projects in Watershed #9 (Middle Missouri River).

The proposed channel restoration was completed over two construction seasons (1998 & 1999), providing a newly created meandering channel with numerous pool, riffle, and run sections. The project incorporated the use of root wads, boulders, footer logs, sod mats, willow clumps and cuttings, coir fabric and seeding of both upland and wetland areas. Sections of floodplain were lowered 1-2 feet to provide areas for wetland development.

According to baseline wetland delineation maps (Barnum and Hoffer 1997) and aerial photographs provided in the environmental assessment prepared for the project by FWP, approximately 7.86 acres of shrub/scrub and emergent wetland occurred within the current monitoring area prior to project implementation (note: reference to a FWS/NRCS delineation resulting in over 14 acres of pre-existing wetlands was found in the project files, but no evidence of such a delineation was found in MDT, NRCS, or FWP project files, and pre-project aerial photographs do not support a 14-acre delineation within the current monitoring area). Hydrology for many of the existing wetlands was thought to be provided by leaking water pipes, with little or no connection to the incised Big Spring Creek channel. The proposed stream restoration was intended to create approximately 1.5 acres of additional wetland habitat, and restore and enhance existing wetlands by reconnecting them with Big Spring Creek.

Target wetland communities to be produced at the site included shallow marsh/wet meadow and wet meadow/scrub-shrub (Inter-Fluve, Inc. 1998). Target wetland functions to be provided at the site included habitat diversity, flood control & storage, threatened/endangered species habitat, general wildlife habitat, sediment filtration, shoreline stabilization, food chain support, nutrient cycling, and uniqueness (Inter-Fluve, Inc. 1998).

FIGURE 1. PROJECT LOCATION
Big Spring Creek
Mitigation Site



PROJECT #: 130091.029
 DATE: MAY 2001
 LOCATION:
 PROJECT MANAGER: B. DUTTON
 DRAWN BY: B. NOECKER



1120 CEDAR PO BOX 8254 MISSOULA, MT 59807

As originally proposed by FWP, the newly created channel was not immediately activated following construction, but was given approximately one year to establish streamside vegetation for stabilization purposes. Water was turned into the new channel in the fall of 2000. This site was first monitored in 2001, and is scheduled to be monitored two times per year over the contract period to document wetland and other biological attributes. The monitoring area is illustrated in **Figure 2 (Appendix A)**.

No performance standards or success criteria were required by the COE or other agencies. The COE determined that the maximum allowable credit at the site is 7.21 acres (Rabbe 1998). This conclusion was subjectively based on acreages of existing and developed wetlands, changes in functions and values, re-creation of a functioning floodplain, and modifications to supporting hydrology (Rabbe 1998). It was the Corps' opinion that the proposed project, while improving the existing setting, would not result in doubling of actual wetland acreage but could essentially double wetland values while establishing "natural" supporting hydrology for the whole complex (Rabbe 1998).

2.0 METHODS

2.1 Monitoring Dates and Activities

The site was visited on June 3rd (spring) and August 5th (mid-season) 2004. The primary purpose of the spring visit was to conduct a bird/general wildlife reconnaissance. The late-May to early-June period was selected for the spring visit because monitoring between mid-May and early June is likely to detect migrant as well as early nesting activities for a variety of avian species (Carlson pers. comm.), as well as maximizing the potential for amphibian detection. In Montana, most amphibian larval stages are present by early June (Werner pers. comm.).

The mid-season visit was conducted to document vegetation, soil, and hydrologic conditions used to map jurisdictional wetlands. All information contained on the Wetland Mitigation Site Monitoring Form (**Appendix B**) was collected at this time. Activities and information conducted/collected included: wetland delineation; wetland/open water boundary mapping; vegetation community mapping; vegetation transects; soils data; hydrology data; bird and general wildlife use; photograph points; macroinvertebrate sampling; functional assessment; and examination of stream habitat conditions including bank stability, fisheries habitat and survival of planted woody vegetation.

2.2 Hydrology

Hydrologic indicators were evaluated at the site during the mid-season visit. Information found in project files indicate that the leaking water pipes on or near the property have been fixed and are no longer contributing to wetland hydrology at the site. The approximate designed channel location is shown on the conceptual restoration plan in **Appendix D**. Wetland hydrology indicators were recorded using procedures outlined in the COE 1987 Wetland Delineation Manual (Environmental Laboratory 1987). Hydrology data were recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**).

All additional hydrologic data were recorded on the mitigation site monitoring form (**Appendix B**). The boundary between wetlands and open water (no rooted vegetation) aquatic habitats was mapped on the aerial photograph and an estimate of the average water depth at this boundary was recorded.

No groundwater monitoring wells were installed at the site. If located within 18 inches of the ground surface (soil pit depth for purposes of delineation), groundwater depths were documented on the routine wetland delineation data form at each data point.

2.3 Vegetation

General dominant species-based vegetation community types (e.g., *Typha latifolia/Scirpus acutus*) were delineated on an aerial photograph during the mid-season visit. Standardized community mapping was not employed as many of these systems are geared towards climax vegetation and may not reflect yearly changes. Estimated percent cover of the dominant species in each community type was listed on the site monitoring form (**Appendix B**).

The 10-foot wide belt transect that was established in 2001 was evaluated for the fourth time **Figure 2 (Appendix A)**. Percent cover was estimated for each vegetative species for each vegetation community encountered within the “belt” using the following values: + (<1%); 1 (1-5%); 2 (6-10%); 3 (11-20%); 4 (21-50%); and 5 (>50%).

The purpose of the transect is to evaluate changes over time, especially the establishment and increase of hydrophytic vegetation. The transect location was marked on the air photo and all data recorded on the mitigation site monitoring form. Transect endpoint locations were recorded with the GPS unit in 2001. Wooden stakes were installed in 2001 to physically mark the transect ends. Photos of the transect were taken from both ends during the mid-season visit.

A comprehensive plant species list for the site was first compiled in 2001 and was updated as new species were encountered. Ultimately, observations from past years will be compared with new data to document vegetation changes over time.

Fourteen woody species were planted at this mitigation site. Planting lists are provided in **Appendix D**. No planting map was available; consequently, not all planting locations were known, and it was not possible for observers to inventory all planted species. Rather, observers recorded the number of dead planted species observed and compared them to known planting numbers.

2.4 Soils

Soils were evaluated during the mid-season visit according to hydric soils determination procedures outlined in the COE 1987 Wetland Delineation Manual. Soil data was recorded for each wetland determination point on the COE Routine Wetland Delineation Data Form (**Appendix B**). The most current terminology used by NRCS was used to describe hydric soils (USDA 1998).

2.5 Wetland Delineation

A wetland delineation of the mitigation site was conducted during the 2001 mid-season visit according to the 1987 COE of Engineers Wetland Delineation Manual. The delineated boundaries were verified and changes made if necessary during the 2002, 2003, and 2004 monitoring. Wetland and upland areas within the monitoring area were investigated for the presence of wetland hydrology, hydrophytic vegetation and hydric soils. The indicator status of vegetation was derived from the National List of Plant Species that occur in Wetlands: Northwest (Region 9) (Reed 1997).

The information was recorded on COE Routine Wetland Delineation Data Forms (**Appendix B**). The wetland/upland boundary was delineated on the air photo and recorded with a resource grade GPS unit in 2001. Minor changes in wetland boundaries were noted in 2004 and drawn onto project maps. These changes were not surveyed with GPS during the 2004 monitoring. The wetland/upland boundary in combination with the wetland/open water habitat boundary was used to calculate the wetland area developed within the monitoring area.

2.6 Mammals, Reptiles, and Amphibians

Mammal, reptile, and amphibian species observations and other positive indicators of use, such as vocalizations, were recorded on the wetland monitoring form during each visit. Indirect use indicators, including tracks; scat; burrows; eggshells; skins; bones; etc., were also recorded. Observations were recorded as the observer traversed the site while conducting other required activities. Direct sampling methods, such as snap traps, live traps, and pitfall traps, were not implemented. A comprehensive list of observed species was compiled. Observations from past years will ultimately be compared with new data.

2.7 Birds

Bird observations were recorded during each visit. No formal census plots, spot mapping, point counts, or strip transects were conducted. During the spring visit, observations were recorded in compliance with the bird survey protocol in **Appendix E**. During the mid-season visit, bird observations were recorded incidental to other monitoring activities. During both visits, observations were categorized by species, activity code, and general habitat association (see data forms in **Appendix B**). Observations from past years will be compared with new data.

2.8 Macroinvertebrates

One macroinvertebrate sample was collected during the mid-season site visit and data recorded on the wetland mitigation monitoring form. Macroinvertebrate sampling procedures and analysis are included in **Appendix F**. The approximate location of this sample point, within emergent marsh habitat in the north portion of the site, is shown on **Figure 2, Appendix A**. The sample was preserved as outlined in the sampling procedure and sent to a laboratory for analysis.

2.9 Functional Assessment

Functional assessment forms were completed for various assessment areas within the monitoring area using the 1999 MDT Montana Wetland Assessment Method. Field data necessary for this assessment were generally collected during the mid-season site visit. The remainder of the functional assessment was completed in the office.

The pre-project functional assessment of the mitigation site was completed using the 1997 MDT wetland assessment method. Thus, while pre- and post-project functional assessment results are not directly comparable, general trends can be discussed.

2.10 Photographs

Photographs were taken during the mid-season visit showing the current land use surrounding the site, the upland buffer, the monitored area, macroinvertebrate sampling location, and the vegetation transect. Each photograph point location was recorded with a resource grade GPS during the 2001 monitoring. The approximate location of photo points is shown on **Figure 2, Appendix A**. All photographs were taken using a 50 mm lens. A description and compass direction for each photograph was recorded on the wetland monitoring form.

2.11 GPS Data

During the 2001 monitoring season, data were collected with a resource grade GPS unit at the vegetation transect beginning and ending locations, at all photograph locations, and at the macroinvertebrate sampling location. Wetland boundaries were also mapped with a resource grade GPS unit. No new GPS data were collected in 2004.

2.12 Maintenance Needs

The newly constructed channel was examined for signs of erosion and channel migration. Where encountered, current or future potential problems were documented, photographed and conveyed to MDT.

3.0 RESULTS

3.1 Hydrology

According to the Western Regional Climate Center, Lewistown yearly precipitation totals for 2001 (12.37 inches), 2002 (15.94 inches), 2003 (13.86), and 2004 (11.96) were 68, 87, 76, and 65 percent, respectively, of the total annual mean precipitation (18.30 inches) in this area.

Inundation was present, to some extent, at all wetlands within the monitoring area during the mid-season visit despite the sub-normal precipitation year. Big Spring Creek contained the only “open water” on the site. Water depths at open water/rooted vegetation interfaces along the

creek ranged between approximately one to two feet. Open water areas are shown on **Figure 3 (Appendix A)**. Specific recorded values are provided on the attached data forms.

Overall, the site was approximately 40 percent inundated, with an average depth of two to four inches and a range of depths from 0 to an estimated four feet. Deepest areas were located at stream pools.

A groundwater component contributes strongly to this site, likely resulting at least partially from alluvial flow. Groundwater was encountered within about 1 foot of the ground surface at most wetlands. Several groundwater discharge sites occur along the toe of the highway fill between the parking area and the northeast corner of the monitoring area. This area is developing very strong wetland characteristics despite early attempts to drain this area with small hand dug ditches. According to MDT, wetlands are not necessarily desirable in this area, as they may be in conflict with future highway expansion (Urban pers. comm.).

3.2 Vegetation

Vegetation species identified on the site are presented in **Table 1** and on the attached data form. No new species were encountered during the 2004 monitoring. Three primary wetland community types were identified and mapped on the mitigation area (**Figure 3, Appendix A**). These included Type 1: *Agrostis alba*, Type 2: *Typha latifolia*, and Type 3: *Salix*. Dominant species within each of these communities are listed on the attached data form (**Appendix B**). Type 1 occurs commonly and intermittently as narrow fringes along the immediate stream channel. Type 2 occurs within emergent marsh communities throughout the site, and Type 3 occurs primarily in association with streamside areas in the south portion of the site.

Upland communities are primarily dominated by seeded and/or weedy herbaceous species including quackgrass (*Agropyron repens*), bearded wheatgrass (*Agropyron caninum*), intermediate wheatgrass (*Agropyron intermedium*), sowthistle (*Sonchus arvensis*), ragweed (*Ambrosia trifida*), field pennycress (*Thlaspi arvense*) and white sweetclover (*Melilotus alba*). A large “transitional upland” area first identified in 2001 occurs west of the creek, and south of the parking lot. This area continues to exhibit signs of transitioning from upland to wetland (**Figure 2 in Appendix A**). Transitional upland areas identified in 2002 and 2003 in the old creek location parallel to the highway and both north and south of the parking area continue to exhibit signs of transitioning from upland to wetland.

Table 1: 2001 - 2004 Big Spring Creek vegetation species list.

Scientific Name	Region 9 (Northwest) Wetland Indicator
<i>Achillea millefolium</i>	FACU
<i>Agropyron caninum</i>	FAC-
<i>Agropyron intermedium</i>	--
<i>Agropyron repens</i>	FACU
<i>Agrostis alba</i>	FACW
<i>Alopecurus pratensis</i>	FACW
<i>Ambrosia trifida</i>	--
<i>Arctium minus</i>	--
<i>Aster spp.</i>	--
<i>Avena fatua</i>	--
<i>Beckmannia syzigachne</i>	OBL
<i>Betula occidentalis</i>	FACW
<i>Bidens cernua</i>	FACW+
<i>Bromus inermis</i>	--
<i>Calamagrostis inexpansa</i>	FACW
<i>Carex aquatilis</i>	OBL
<i>Carex nebrascensis</i>	OBL
<i>Carex utriculata</i>	OBL
<i>Cirsium arvense</i>	FAC-
<i>Cornus stolonifera</i>	FACW
<i>Crataegus douglasii</i>	FAC
<i>Dactylis glomerata</i>	--
<i>Echinochloa crusgalli</i>	FACW
<i>Eleocharis palustris</i>	OBL
<i>Elodea canadensis</i>	OBL
<i>Epilobium ciliatum</i>	FACW-
<i>Equisetum arvense</i>	FAC
<i>Fraxinus pensylvanica</i>	FAC
<i>Galium aparine</i>	--
<i>Glyceria elata</i>	FACW+
<i>Glycyrrhiza lepidota</i>	FAC+
<i>Hordeum jubatum</i>	FAC-
<i>Iva xanthifolia</i>	FAC
<i>Juncus bufonius</i>	FACW+
<i>Juncus ensifolius</i>	FACW
<i>Juncus nodosus</i>	OBL
<i>Juncus torreyi</i>	FACW
<i>Lactuca serriola</i>	FACU
<i>Lemna minor</i>	OBL
<i>Linaria vulgaris</i>	--
<i>Lycopus americanus</i>	OBL
<i>Medicago lupulina</i>	FAC
<i>Melilotus alba</i>	FACU
<i>Melilotus officinalis</i>	FACU
<i>Mentha arvensis</i>	FAC
<i>Muhlenbergia minutissima</i>	FAC
<i>Nasturtium officinale</i>	OBL
<i>Phalaris arundinacea</i>	FACW
<i>Phleum pratense</i>	FAC-
<i>Plantago major</i>	FAC+
<i>Poa pratensis</i>	FAC

Table 1 (continued): 2001 - 2004 Big Spring Creek vegetation species list.

Scientific Name	Region 9 (Northwest) Wetland Indicator
<i>Polygonum lapathifolium</i>	FACW
<i>Polypogon monspeliensis</i>	FACW
<i>Populus angustifolia</i>	FACW
<i>Populus deltoides</i>	FAC
<i>Populus tremuloides</i>	FAC+
<i>Populus trichocarpa</i>	FAC
<i>Prunus virginiana</i>	FACU
<i>Ribes aureum</i>	FAC+
<i>Ranunculus aquatilis</i>	OBL
<i>Rosa woodsii</i>	FACU
<i>Rumex crispus</i>	FACW
<i>Sagittaria cuneata</i>	OBL
<i>Salix amygdaloides</i>	FACW
<i>Salix exigua</i>	OBL
<i>Salix lutea</i>	OBL
<i>Scirpus acutus</i>	OBL
<i>Scirpus microcarpus</i>	OBL
<i>Scirpus pungens</i>	OBL
<i>Shepherdia canadensis</i>	--
<i>Sisymbrium altissimum</i>	FACU-
<i>Sium suave</i>	OBL
<i>Solidago canadensis</i>	FACU
<i>Sonchus arvensis</i>	FACU+
<i>Taraxacum officinale</i>	FACU
<i>Thlaspi arvense</i>	--
<i>Trifolium fragiferum</i>	FACU
<i>Trifolium repens</i>	--
<i>Typha latifolia</i>	OBL
<i>Verbascum thapsus</i>	--

Vegetation transect results in 2004 differed from the 2003 results near the middle and east end of the transect, where Type 2 habitat continues to encroach into previously identified upland areas. Vegetation transect results are detailed in the attached data form, and are summarized in **Charts 1 and 2** and **Table 2**.

Chart 1: Transect maps showing vegetation types from the start of transect (0 feet) to the end of transect (418 feet) for each year monitored.





Table 2: Vegetation transect data summary.

Monitoring Year	2001	2002	2003	2004
Transect Length (feet)	418	418	418	418
# Vegetation Community Transitions along Transect	8	8	8	4
# Vegetation Communities along Transect	3	3	3	3
# Hydrophytic Vegetation Communities along Transect	2	2	2	2
Total Vegetative Species	31	31	31	31
Total Hydrophytic Species	23	23	23	23
Total Upland Species	8	8	8	8
Estimated % Total Vegetative Cover	95	95	95	95
% Transect Length Comprised of Hydrophytic Vegetation Communities	71	73	89	99
% Transect Length Comprised of Upland Vegetation Communities	29	27	11	1
% Transect Length Comprised of Unvegetated Open Water	0	0	0	0
% Transect Length Comprised of Bare Substrate	0	0	0	0

Chart 2: Length of vegetation communities along Transect 1.



Numerous willow cuttings and other woody species were planted as part of the overall revegetation plan for the project. Additionally, the NRCS and American Foresters Society sponsored a community project at the site that resulted in additional plantings. Observed mortality of planted woody vegetation species is summarized below in **Table 3**. As specific planting locations were unknown, only observations of dead, obviously planted individuals were recorded in order to avoid spending available monitoring time searching the site for possible planting areas.

Table 3: 2004 observed mortality of planted woody species.

Species	Estimated # Originally Planted	Observed # Dead	Comments
<i>Salix exigua</i> <i>Salix amygdaloides</i>	up to 3,500 cuttings; species not distinguished	see comments	Willows planted below the ordinary high water mark were generally dead, presumably due to drowning. Willows planted above the OHWM were generally alive. Estimated overall survival rate of 50 – 60%.
<i>Populus deltoides</i>	21	10	Mortality likely due to drier or wetter than anticipated conditions at individual planting locations.
<i>Populus trichocarpa</i>	24	11	Mortality likely due to drier or wetter than anticipated conditions at individual planting locations.
<i>Populus angustifolia</i>	30	0	Doing well; many observed.
<i>Populus tremuloides</i>	50	0	No dead observed, but estimated <50 live observed. Assume some mortality.
<i>Betula occidentalis</i>	31	5	Few dead observed, but estimated <10 live observed. Mortality likely due to drought.
<i>Rosa woodsii</i>	10	0	No dead observed, but estimated <5 live observed. Mortality likely due to drought / competition with upland grasses.
<i>Cornus stolonifera</i>	130	0	No dead observed, but estimated <50 live observed. Mortality likely due to drought / competition with upland grasses, and possibly deer.
<i>Prunus virginiana</i>	150	10	Doing well; numerous observations.
<i>Shepherdia canadensis</i>	30	0	No dead observed, but estimated <20 live observed. Assume some mortality.
<i>Fraxinus pennsylvanica</i>	30	0	Doing well; several observed.
<i>Ribes aureum</i>	35	0	No dead observed, but estimated <10 live observed.
<i>Crataegus douglasii</i>	10	2	Few live or dead observed.

3.3 Soils

According to the Fergus County soil survey (Soil Conservation Service 1988), pre-existing soils at the site were mapped as Fluvaquentic Haplaquolls and Enbar-Nesda loams. Fluvaquentic Haplaquolls are poorly drained soils on flood plains that formed in alluvium. Enbar-Nesda loams are well drained to somewhat poorly drained soils that occur on floodplains and terraces. Oddly, soils descriptions provided in the survey for these two map units seem to apply in the reverse on the ground. The survey describes the upland portions of the site as supporting the wetter Fluvaquentic Haplaquolls, and the wetland portions as supporting drier Enbar-Nesda loams. On the ground, just the opposite seems true. Both of these soils types exhibit a seasonal high water table. Fluvaquentic Haplaquolls are included on the Fergus County hydric soils list (floodplains), while Enbar-Nesda loams are not considered hydric.

Soils sampled in wetland areas were generally comprised of silty clay loams or silt loams with a matrix color of 10YR3/1 without mottles, or 10YR3/2 with distinct mottles in the range of 10YR 4/6, indicating a fluctuating water table. Wetland soils were saturated or inundated at the time of the survey.

3.4 Wetland Delineation

Delineated wetland boundaries are illustrated on **Figure 3**. Completed wetland delineation forms are included in **Appendix B**. Soils, vegetation, and hydrology are discussed in preceding sections. Total wetland area for the site increased by 0.73 acres in 2004 as shown on **Figure 3**. Delineation results including the expanded areas are as follows:

Big Spring Creek: 10.44 wetland acres
2.41 acres open water (non-wetland perennial stream channel)

Based on maps provided in the project EA, approximately 7.86 wetland acres and 1.3 acres of non-wetland perennial stream channel occurred within the monitoring area prior to project implementation. Currently, the site has gained 2.58 wetland acres and 1.11 acres of non-wetland perennial stream channel.

3.5 Wildlife and Fish

Wildlife species, or evidence of wildlife, observed on the site during 2004 monitoring efforts are listed in **Table 4** in bold, with the remaining listed species having been seen during previous years monitoring. Specific evidence observed, as well as activity codes pertaining to birds, are provided on the completed monitoring form in **Appendix B**. Six mammal, one reptile, one amphibian, and 26 bird species were noted using portions of the mitigation site during 2004 monitoring. Rainbow trout (*Oncorhynchus mykiss*) were also observed. The wetland and stream habitat provided on the site, particularly large streamside wetland complexes in the north and south portions of the site, provide quality wildlife habitat for several species. This habitat value is expected to increase as vegetation establishes and diversifies, and as additional wetlands are restored/created. The lone wood duck nesting box located on the site (see **Figure 2, Appendix A**) appeared to be inactive during the 2003 and 2004 nesting seasons.

Preliminary fish shocking data for the restored reach are encouraging. In 2001, the reach of Big Spring Creek including the restored channel was shocked, and yielded 710 rainbow and brown (*Salmo trutta*) trout over 10 inches in length (MFWP 2002). This compares with pre-project (1995 – 2000) shocking results that averaged 434 trout over 10 inches in length (MFWP 2002) through reaches including the project area.

3.6 Macroinvertebrates

Macroinvertebrates were sampled within the emergent marsh complex east of the creek in the north portion of the site (see **Figure 2**). The same location was sampled during each of the four monitoring seasons (**Chart 3**). Macroinvertebrate sampling results are provided in **Appendix F** and were summarized by Rhithron Associates in the italicized sections below (Bollman 2004).

Scores indicate that sub-optimal conditions existed in the initial three years of sampling at the Big Spring Creek site (Chart 3). In 2004, scores suggest that conditions improved to optimal (Chart 3). Taxa richness increased dramatically in 2004, more than doubling the previous year's sampling. Macrophytes apparently contributed to habitat diversity here. The mayfly Callibaetis spp., which was absent from the collection of 2002, and which reappeared at the site in 2003, maintained its importance to the fauna in 2004. The invertebrate assemblage gave evidence of colonization of all important habitats; water column, macrophytes, and benthic substrates. Moderate elevation of the biotic index value between 2003 and 2004 may have been related to warmer water temperatures in the latter year.

Chart 3: Macroinvertebrate bioassessment scores 2001 – 2004.




Table 4: Fish and wildlife species observed on the Big Spring Creek Mitigation Site 2001-2004.

FISH	
Rainbow Trout (<i>Oncorhynchus mykiss</i>)	
AMPHIBIANS	
Western Chorus Frog (<i>Pseudacris triseriata</i>)	
REPTILES	
Western Terrestrial Garter Snake (<i>Thamnophis elegans</i>)	
BIRDS	
American Robin (<i>Turdus migratorius</i>) Black-headed Grosbeak (<i>Pheucticus melanocephalus</i>) Belted Kingfisher (<i>Ceryle alcyon</i>) Black-billed Magpie (<i>Pica pica</i>) Blue-winged Teal (<i>Anas discors</i>) Canada Goose (<i>Branta Canadensis</i>) Cedar Waxwing (<i>Bombycilla cedrorum</i>) Cinnamon Teal (<i>Anas cyanoptera</i>) Cliff Swallow (<i>Petrochelidon pyrrhonota</i>) Common Merganser (<i>Mergus merganser</i>) Common Snipe (<i>Gallinago gallinago</i>) Common Yellowthroat (<i>Geothlypis trichas</i>) Downy Woodpecker (<i>Picoides pubescens</i>) Eastern Kingbird (<i>Tyrannus tyrannus</i>) European Starling (<i>Sturnus vulgaris</i>) Gray Catbird (<i>Dumetella carolinensis</i>) Killdeer (<i>Charadrius vociferous</i>) Mallard (<i>Anas platyrhynchos</i>)	Mourning Dove (<i>Zenaida macroura</i>) Northern Flicker (<i>Colaptes auratus</i>) Northern Harrier (<i>Circus cyaneus</i>) Northern Rough-winged Swallow (<i>Stelgidopteryx serripennis</i>) Osprey (<i>Pandion haliaetus</i>) Red-tailed Hawk (<i>Buteo jamaicensis</i>) Red-winged Blackbird (<i>Agelaius phoeniceus</i>) Ring-necked Pheasant (<i>Phasianus colchicus</i>) Sandhill Crane (<i>Grus Canadensis</i>) Song Sparrow (<i>Melospiza melodia</i>) Sora (<i>Porzana Carolina</i>) Spotted Sandpiper (<i>Actitis macularia</i>) Tree Swallow (<i>Tachycineta bicolor</i>) Willow Flycatcher (<i>Empidonax traillii</i>) Yellow Warbler (<i>Dendroica petechia</i>) Yellow-headed Blackbird (<i>Xanthocephalus xanthocephalus</i>) Yellow-rumped Warbler (<i>Dendroica coronata</i>)
MAMMALS	
White-tailed Deer (<i>Odocoileus virginianus</i>) American Beaver (<i>Castor Canadensis</i>) Eastern Cottontail (<i>Sylvilagus floridanus</i>) Meadow Vole (<i>Microtus pennsylvanicus</i>) Muskrat (<i>Ondatra zibethicus</i>) Raccoon (<i>Procyon lotor</i>)	

Bolded species were seen during the 2004 monitoring. All other species have been seen during one or more of the previous monitoring seasons.

3.7 Functional Assessment

Completed functional assessment forms are presented in **Appendix B**. Functional assessment results in 2004 were virtually unchanged from the 2001 - 2003 assessments, and are summarized in **Table 5**. For comparative purposes, the functional assessment results for baseline conditions prepared by Inter-Fluve are also included in **Table 5**. However, the baseline assessment was performed using a modified 1997 MDT assessment method. Several parameters of this method were substantially revised during development of the 1999 MDT assessment method, which was applied during 2004 monitoring. For example, baseline fish habitat scored a 1.0 using the 1997

method, and scored a 0.9 post project using the 1999 method due to the addition of several variables for consideration in the updated method. Fish habitat increased dramatically with addition of channel length, substrate improvement, and other features; however, this was not reflected in the comparative functional assessments. Thus, direct comparison of pre- and post-project functions is not possible, although some general trends can be noted. Also, as the baseline assessment was performed using a modified 1997 MDT method, it resulted in an incorrect overall category designation (Category IV). This was corrected to a Category III on **Table 5**.

Large wetland polygons bisected by the stream rated as Category II sites, primarily due to high wildlife and fish habitat, flood attenuation, sediment removal, production export, and recreation/education ratings. Narrow fringes along the creek rated as Category III sites, rating high for groundwater discharge and recreation/education. Isolated depressions rated as Category III sites and scored high for sediment/nutrient removal and groundwater discharge.

Generally speaking, functions that increased substantially over baseline conditions include wildlife and fish habitat, flood attenuation, sediment/nutrient/toxicant removal, production export, and groundwater discharge. The pre-project site provided about 29 functional units within the monitoring area (using the 1997 method), and the post-project site provides about 90 functional units (using the 1999 method), for a conservative gain of at least 61 functional units.

3.8 Photographs

Representative photographs taken from photo-points and transect ends are provided in **Appendix C**. A 2004 aerial photograph is also provided in **Appendix C**.

3.9 Maintenance Needs/Recommendations

Although a thorough investigation of all stream banks was not completed, it does appear that the outside bend of the creek immediately south of the designated parking area is experiencing some minor lateral migration. The one Wood Duck box on the site was hanging upside down on the tree that it is attached to. This problem should be corrected to encourage use of the box by cavity nesting species.

Table 5: Summary of 2004 wetland function/value ratings and functional points¹ at the Big Spring Creek Mitigation Project.

Function and Value Parameters From the 1999 MDT Montana Wetland Assessment Method	Wetland Sites			
	2004: Large wetland polygons bisected by creek near north, east, and south ends of site.	2004: Isolated wetland depressions west of creek.	2004: Narrow wetland fringe segments along creek.	1998: Baseline assessment. ²
Listed/Proposed T&E Species Habitat	Low (0.3)	Low (0.0)	Low (0.3)	Low (0.2)
MNHP Species Habitat	Mod (0.6)	Low (0.1)	Low (0.1)	Low (0.0)
General Wildlife Habitat	High (0.9)	Mod (0.5)	Mod (0.7)	Mod (0.5)
General Fish/Aquatic Habitat	High (0.9)	NA	Mod (0.7)	High (1.0)
Flood Attenuation	High (0.7)	Low (0.2)	Low (0.2)	Low (0.3)
Short and Long Term Surface Water Storage	Mod (0.6)	Low (0.3)	Low (0.3)	--
Sediment, Nutrient, Toxicant Removal	High (1.0)	High (1.0)	Mod (0.6)	Low (0.1)
Sediment/Shoreline Stabilization	Mod (0.7)	NA	Mod (0.7)	NA
Production Export/Food Chain Support	High (0.9)	Low (0.3)	Mod (0.4)	Low (0.4)
Groundwater Discharge/Recharge	High (1.0)	High (1.0)	High (1.0)	NA
Uniqueness	Low (0.3)	Low (0.3)	Low (0.3)	Low (0.2)
Recreation/Education Potential	High (1.0)	Mod (0.5)	High (1.0)	High (1.0)
Actual Points/Possible Points	8.9 / 12	4.2 / 10	5.3 / 12	3.7 / 10
% of Possible Score Achieved	74%	42%	44%	37%
Overall Category	II	III	III	III ³
Total Acreage of Assessed Wetlands within AA Boundaries (note: non-wetland stream channel is not included in these totals) * Pre-project (baseline) wetland areas within the current monitoring area boundaries were measured via digital planimeter from delineation maps provided in project EA.	9.84	0.54	0.06	7.86
Functional Units (acreage x actual points)	87.6	2.3	0.3	29.1
Net Acreage Gain	Site currently supports 10.44 wetland acres and 2.4 non-wetland perennial stream channel acres. Baseline conditions within the current monitoring area boundaries included 7.86 wetland acres and 1.3 acres of non-wetland perennial stream channel. Net gain is approximately 2.58 wetland acres and 1.1 acres of non-wetland perennial stream channel.			
Net Functional Unit Gain²	Approximately 61.1 Functional Units ²			

¹ See completed MDT functional assessment forms in Appendix B for further detail.

² The baseline assessment was performed by Inter-Fluve using a modified 1997 MDT assessment method. Several parameters were substantially revised and applied to the 1999 MDT assessment method, which was applied during 2004 monitoring. Thus, direct comparison of pre- and post-project functions are not possible, but general trends can be noted.

³ The baseline assessment was performed using a modified 1997 MDT method, which resulted in an incorrect overall category designation (Category IV). This was corrected to a Category III.

3.10 Current Credit Summary

Approximately 10.44 wetland acres and 2.4 acres of non-wetland perennial stream channel occur within the monitoring area. Based on maps provided in the project EA, approximately 7.86 wetland acres and 1.3 acres of non-wetland perennial stream channel occurred within the monitoring area prior to project implementation. Currently, the site has gained 2.58 wetland acres and 1.11 acres of non-wetland perennial stream channel, substantially improving fish habitat. It was originally anticipated that the area encompassed by the old stream channel would develop into upland riparian habitat following construction; however, this area continues to transition to emergent marsh and scrub/shrub wetland thus providing wetland mitigation acreage that was not originally anticipated.

The pre-project site provided about 29 functional units within the monitoring area (using the 1997 method), and the post-project site provides about 90 functional units (using the 1999 method), for a conservative gain of at least 61 functional units.

The COE determined that the maximum allowable credit at the site is 7.21 acres (Rabbe 1998). This conclusion was subjectively based on acreages of existing and developed wetlands, changes in functions and values, re-creation of a functioning floodplain, and modifications to supporting hydrology (Rabbe 1998). No performance standards were required by the COE, although the site appears to be well on its way to functioning as anticipated.

4.0 REFERENCES


- Bollman, W. 2004. MDT Wetland Mitigation Monitoring Project – Aquatic Invertebrate Monitoring Summary 2001, 2002, 2003, 2004. Rhithron Associates Inc. Missoula, MT.
- Carlson, J. Program Zoologist, MT Natural Heritage Program. Helena, MT. April 2001 comm.
- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. US Army Corps of Engineers. Washington, DC.
- Montana Fish, Wildlife & Parks. 2002. Preliminary Brewery Flats fish shocking data. Lewistown, MT. Unpublished data.
- Rabbe, M. 1998. June 2, 1998 letter from Michael Rabbe (U.S. Army Corps of Engineers) to Michael Rotor (Inter-Fluve, Inc.) regarding crediting on the Big Spring Creek project – Corps File # 199790594. Helena, MT.
- Ralph, C.J., Geupel, G.R., Pyle, P., Martin, T.E., and D.F. DeSante. 1993. *Handbook of field methods for monitoring landbirds*. Gen. Tech. Rep. PSW-GTR-144. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Dept. of Agriculture. 41 p.
- Reed, P.B. 1988. National list of plant species that occur in wetlands: North West (Region 9). Biological Report 88(26.9), May 1988. U.S. Fish & Wildlife Service. Washington, D.C.
- Rotar, M. Water Resources Engineer, Inter-Fluve, Inc. Bozeman, MT. May 28, 1998 letter to Mr. Doug McDonald, U.S. Army Corps of Engineers. Helena, MT.
- Rotar, M. Water Resources Engineer, Inter-Fluve, Inc. Bozeman, MT. May, 1998 Montana Wetland Field Evaluation Form (rev. 9/23/1997, Draft) for Big Spring Creek Renaturalization Project.
- Soil Conservation Service. 1988. Soil survey of Fergus County, Montana. Bozeman, MT.
- Urban, L. Wetland Mitigation Specialist, Montana Department of Transportation. Helena, MT. March 2001 meeting, May 2001 & March 2002 telephone conversations.
- USDA Natural Resources Conservation Service. 1998. *Field Indicators of Hydric Soils in the United States*, Version 4. G. Hurt, P. Whited and R. Pringle (eds.). USDA, NRCS Fort Worth, TX.
- Werner, K. Herpetologist, Salish-Kootenai Community College. Pablo, MT. May 1998 instructional presentation.

Appendix A

FIGURES 2 & 3

*MDT Wetland Mitigation Monitoring
Big Spring Creek
Lewistown, Montana*

Figure 2 – Monitoring Activity Locations




Legend

- Monitoring Area Limits
- Photograph Points
- Aerial Reference Points
- Macro-invertebrate Sample Points
- Vegetation Transect
- Bird Box
- Base Photograph Date: July 11, 2001

SCALE 1" = 200 ft

PROJECT NAME MDT Big Spring Creek Wetland Mitigation	
DRAWING TITLE Monitoring Activity Locations	
PROJ NO: 330054.502	DRAWN: RA
FILE NAME: TASK502BASE.dwg	CHECKED:
SCALE: 1" = 200ft	APPVD: BD
LOCATION: Big Spring Creek	PROJ MGR: BD
SHEET NUMBER 2 OF	
REV - DATE: 3-23-05	

Figure 3 – Mapped Site Features 2004



Appendix B

**COMPLETED 2004 WETLAND MITIGATION SITE MONITORING
FORM**

COMPLETED 2004 BIRD SURVEY FORMS

COMPLETED 2004 WETLAND DELINEATION FORMS

COMPLETED 2004 FUNCTIONAL ASSESSMENT FORMS

*MDT Wetland Mitigation Monitoring
Big Spring Creek
Lewistown, Montana*

LWC / MDT WETLAND MITIGATION SITE MONITORING FORM

Project Name: **Big Spring Creek** Project Number: _____ Assessment Date: **8/5/04**
 Location: **Lewistown** MDT District: **Billings** Milepost: _____
 Legal description: **T15N R18E** Section **23** Time of Day: **0800-1100**
 Weather Conditions: **Partly cloudy & warm approx. 70 degrees** Person(s) conducting the assessment:
Traxler
 Initial Evaluation Date: **8 / 29 / 01** Visit #: **2** Monitoring Year: **2004 (year 4)**
 Size of evaluation area: **15 acres** Land use surrounding wetland: **Park, Residential, industrial**

HYDROLOGY

Surface Water Source: **Big Spring Creek, groundwater**
 Inundation: Present Absent _____ Average depths: **.25ft** Range of depths: **0 - 4 ft**
 Assessment area under inundation: **40%**
 Depth at emergent vegetation-open water boundary: **1-2 ft**
 If assessment area is not inundated are the soils saturated w/in 12" of surface: Yes No _____
 Other evidence of hydrology on site (drift lines, erosion, stained vegetation etc.): **Most of the wetlands on site were either inundated or saturated to the surface. Spring flow from east side of highway is influencing wetland development in the northeast corner of the site.**

Groundwater

Monitoring wells: Present _____ Absent
 Record depth of water below ground surface

Well #	Depth	Well #	Depth	Well #	Depth

Additional Activities Checklist:

- Map emergent vegetation-open water boundary on air photo
- Observe extent of surface water during each site visit and look for evidence of past surface water elevations (drift lines, erosion, vegetation staining etc.)
- NA GPS survey groundwater monitoring wells locations if present

COMMENTS/PROBLEMS: Area adjacent to toe of road fill north and south of the main parking area is inundated and developing strong wetland characteristics. These areas are groundwater driven and also receive surface flows from springs to the east of the highway.

VEGETATION COMMUNITIES

Community No.: 1 Community Title (main species): AGR ALB

Dominant Species	% Cover	Dominant Species	% Cover
AGR ALB	>50	CAR AQU	11-20
MEN ARV	11-20		
BID CER	1-5		
EQU ARV	11-20		
JUN NOD	11-20		

COMMENTS/PROBLEMS:

Community No.: 2 Community Title (main species): TYP LAT

Dominant Species	% Cover	Dominant Species	% Cover
TYP LAT	>50	SCI PUN	1-5
SCI ACU	6-10	CAR NEB	6-10
AGR ALB	6-10	CAR AQU	6-10
ALO PRA	6-10		
PHA ARU	11-20		

COMMENTS/PROBLEMS: _____

Community No.: 3 Community Title (main species): SALIX

Dominant Species	% Cover	Dominant Species	% Cover
SAL LUT	>50	AGR ALB	6-10
SAL AMY	21-50		
SAL EXI	21-50		
CAL INE	6-10		
MEN ARV	6-10		

COMMENTS/PROBLEMS: Similar to 2003.

Additional Activities Checklist:

Record and map vegetative communities on air photo

COMPREHENSIVE VEGETATION LIST

Species	Vegetation Community Number(s)	Species	Vegetation Community Number(s)
<i>Achillea millefolium</i>	4	<i>Lycopus americanus</i>	1,2
<i>Agropyron caninum</i>	4	<i>Medicago lupulina</i>	4,5
<i>Agropyron intermedium</i>	4	<i>Melilotus alba</i>	4,5
<i>Agropyron repens</i>	4	<i>Melilotus officinalis</i>	4
<i>Agrostis alba</i>	1,2,3,5	<i>Mentha arvensis</i>	1,3,5
<i>Alopecurus pratensis</i>	2,5	<i>Muhlenbergia minutissima</i>	4
<i>Ambrosia trifida</i>	4	<i>Nasturtium officinale</i>	1,2
<i>Arctium minus</i>	4,5	<i>Phalaris arundinacea</i>	1,2,3
<i>Aster spp.</i>	4	<i>Phleum pratense</i>	4
<i>Avena fatua</i>	4	<i>Plantago major</i>	4
<i>Beckmannia syzigachne</i>	1	<i>Poa pratensis</i>	4
<i>Betula occidentalis</i>	3	<i>Polygonum lapathifolium</i>	5
<i>Bidens cernua</i>	1,5	<i>Polypogon monspeliensis</i>	5
<i>Bromus inermis</i>	4	<i>Populus angustifolia</i>	3,4
<i>Calamagrostis inexpansa</i>	1,3	<i>Populus deltoides</i>	3,4
<i>Carex aquatilis</i>	1,2	<i>Populus tremuloides</i>	3,4
<i>Carex nebrascensis</i>	2	<i>Populus trichocarpa</i>	3,4
<i>Carex utriculata</i>	1,2	<i>Prunus virginiana</i>	3
<i>Cirsium arvense</i>	4	<i>Ribes aureum</i>	4
<i>Cornus stolonifera</i>	3	<i>Ranunculus aquatilis</i>	1,2
<i>Crataegus douglasii</i>	4,5	<i>Rosa woodsii</i>	4
<i>Dactylis glomerata</i>	4	<i>Rumex crispus</i>	1,5
<i>Echinochloa crusgalli</i>	5	<i>Sagittaria cuneata</i>	1,2
<i>Eleocharis palustris</i>	1,2	<i>Salix amygdaloides</i>	3
<i>Elodea canadensis</i>	2	<i>Salix exigua</i>	3
<i>Epilobium ciliatum</i>	1,2	<i>Salix lutea</i>	3
<i>Equisetum arvense</i>	1,5	<i>Scirpus acutus</i>	1,2
<i>Fraxinus pensylvanica</i>	4	<i>Scirpus microcarpus</i>	2
<i>Galium aparine</i>	4,5	<i>Scirpus pungens</i>	1
<i>Glyceria elata</i>	1,5	<i>Shepherdia canadensis</i>	4
<i>Glycyrrhiza lepidota</i>	4,5	<i>Sisymbrium altissimum</i>	4
<i>Hordeum jubatum</i>	1,5	<i>Sium suave</i>	1
<i>Iva xanthifolia</i>	4,5	<i>Solidago canadensis</i>	4,5
<i>Juncus bufonius</i>	1	<i>Sonchus arvensis</i>	4
<i>Juncus ensifolius</i>	1	<i>Taraxacum officinale</i>	4
<i>Juncus nodosus</i>	1,2	<i>Thlaspi arvense</i>	4
<i>Juncus torreyi</i>	1	<i>Trifolium fragiferum</i>	4
<i>Lactuca serriola</i>	4,5	<i>Trifolium repens</i>	4
<i>Lemna minor</i>	1,2	<i>Typha latifolia</i>	2
<i>Linaria vulgaris</i>	4	<i>Verbascum thapsus</i>	4

COMMENTS/PROBLEMS: _____

PHOTOGRAPHS

Using a camera with a 50 mm lenses and color film take photographs of the following permanent reference points listed in the checklist below. Record the direction of the photograph using a compass. (The first time at each site establish a permanent reference point by setting a ½ inch rebar or fencepost extending 2-3' above ground, survey the location with a resource grade GPS and mark the location on the air photo.)

Checklist:

- One photo for each of the 4 cardinal directions surrounding wetland
- At least one photo showing upland use surrounding wetland – if more than one upland use exists, take additional photos
- At least one photo showing buffer surrounding wetland
- One photo from each end of vegetation transect showing transect

Location	Photo Frame #	Photograph Description	Compass Reading
A		See photo sheets and field notes	
B			
C			
D			
E			
F			
G			
H			

COMMENTS/PROBLEMS: _____

GPS SURVEYING

Using a resource grade GPS survey the items on the checklist below. Collect at least 3 location points with the GPS unit set at 5 second recording rate. Record file numbers fore site in designated GPS field notebook

Checklist:

- _____ Jurisdictional wetland boundary
- _____ 4-6 landmarks recognizable on the air photo
- _____ Start and end points of vegetation transect(s)
- _____ Photo reference points
- _____ Groundwater monitoring well locations

COMMENTS/PROBLEMS: ___GPS not used during 2003; minor changes in wetland borders were hand-adjusted using aerial photograph and 2002 delineation.

WETLAND DELINEATION

(Attach Corps of Engineers delineation forms)

At each site conduct the items on the checklist below:

- Delineate wetlands according to the 1987 Army Corps manual.
- Delineate wetland-upland boundary on the air photo
- Survey wetland-upland boundary with a resource grade GPS survey

COMMENTS/PROBLEMS: See attached completed delineation forms.

FUNCTIONAL ASSESSMENT

(Complete and attach full MDT Montana Wetland Assessment Method field forms; also attach abbreviated field forms, if used)

COMMENTS/PROBLEMS: See attached completed functional assessment forms.

MAINTENANCE

Were man-made nesting structures installed at this site? YES NO

If yes, do they need to be repaired? YES NO

If yes, describe problems below and indicate if any actions were taken to remedy the problems.

Were man-made structures build or installed to impound water or control water flow into or out of the wetland?

YES NO

If yes, are the structures working properly and in good working order? YES NO

If no, describe the problems below.

COMMENTS/PROBLEMS:

MDT WETLAND MONITORING – VEGETATION TRANSECT

Site: Big Spring Creek Date: 8/5/04 Examiner: Traxler Transect # 1

Approx. transect length: 418 ft Compass Direction from Start (Upland): 94 degrees

Vegetation type A:		TYP LAT (veg type 2)	
Length of transect in this type:	172	feet	
Species:		Cover:	
TYP LAT		>50	
AGR ALB		1-5	
ELE PAL		>50	
MEN ARV		1-5	
JUN NOD		6-10	
CER DEM		1-5	
SAG CUN		1-5	
CAR NEB		6-10	
ALO PRA		6-10	
LEM MIN		11-20	
CAR AQU		1-5	
Total Vegetative Cover:		100%	

Vegetation type B:		AGR ALB (veg type 1)	
Length of transect in this type:	95	feet	
Species:		Cover:	
AGR ALB (11-20)		6-10	
CAL INE		1-5	
EPI CIL		1-5	
MEN ARV		11-20	
BID CER		1-5	
AGR CAN		1-5	
CON MAC		<1	
RUM CRI		<1	
TYP LAT (1-5)		11-20	
CAR NEB		11-20	
ALO PRA		1-5	
Total Vegetative Cover:		100%	

Vegetation type C:		TYP LAT (veg type 2)	
Length of transect in this type:	149	feet	
Species:		Cover:	
TYP LAT		>50	
ALO PRA		1-5	
AGR ALB		1-5	
EPI CIL		1-5	
JUN NOD		11-20	
JUN TOR		1-5	
GLY ELA		1-5	
ELE PAL		21-50	
RUM CRI		1-5	
CAR NEB		6-10	
Total Vegetative Cover:		100%	

Vegetation type D:		Upland	
Length of transect in this type:	2	feet	
Species:		Cover:	
SON ARV		21-50	
HOR JUB		<1	
AGR INT		11-20	
THL ARV		11-20	
PLA MAJ		1-5	
POL LAP		1-5	
TRI FRA		1-5	
AMB TRI		<1	
CIR ARV		1-5	
MEN ARV		1-5	
Total Vegetative Cover:		100	

**DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)**

Project/Site: Big Spring Creek	Project No: Task 29	Date: 5-Aug-2004
Applicant/Owner: Montana Department of Transportation	County: Fergus	State: Montana
Investigators: Traxler	Plot ID: 1	

Do Normal Circumstances exist on the site? <input checked="" type="radio"/> Yes <input type="radio"/> No	Community ID: EM/SS
Is the site significantly disturbed (Atypical Situation:)? <input type="radio"/> Yes <input checked="" type="radio"/> No	Transect ID: NA
Is the area a potential Problem Area? (If needed, explain on the reverse side)	Field Location: E. of stream, S. portion (pre-exist)

VEGETATION (USFWS Region No. 9)

Dominant Plant Species(Latin/Common)	Stratum	Indicator	Plant Species(Latin/Common)	Stratum	Indicator
<i>Salix amygdaloides</i>	Shrub	FACW	<i>Rumex crispus</i>	Herb	FACW
<i>Willow, Peach-Leaf</i>			<i>Dock, Curly</i>		
<i>Salix exigua</i>	Shrub	OBL	<i>Beckmannia syzigachne</i>	Herb	OBL
<i>Willow, Sandbar</i>			<i>Sloughgrass, American</i>		
<i>Agrostis alba</i>	Herb	FACW	<i>Glycyrrhiza lepidota</i>	Herb	FAC+
<i>Redtop</i>			<i>Licorice, American</i>		
<i>Mentha arvensis</i>	Herb	FAC	<i>Cirsium arvense</i>	Herb	FACU+
<i>Mint, Field</i>			<i>Thistle, Creeping</i>		
<i>Typha latifolia</i>	Herb	OBL	<i>Juncus ensifolius</i>	Herb	FACW
<i>Cattail, Broad-Leaf</i>			<i>Rush, Three-Stamen</i>		
<i>Calamagrostis inexpansa</i>	Herb	FACW			
<i>Small-Reedgrass, Narrow-Spike</i>					

Percent of Dominant Species that are OBL, FACW or FAC: FAC Neutral: 8/9 = 88.89%
(excluding FAC-) 10/11 = 90.91% Numeric Index: 23/11 = 2.09

Remarks:

HYDROLOGY

<p><u>YES</u> Recorded Data(Describe in Remarks): <u>NO</u> Stream, Lake or Tide Gauge <u>YES</u> Aerial Photographs <u>NO</u> Other</p> <p><u>NO</u> No Recorded Data</p> <p>Field Observations</p> <p>Depth of Surface Water: N/A (in.)</p> <p>Depth to Free Water in Pit: = 0 (in.)</p> <p>Depth to Saturated Soil: N/A (in.)</p>	<p>Wetland Hydrology Indicators</p> <p>Primary Indicators</p> <p><u>NO</u> Inundated <u>YES</u> Saturated in Upper 12 Inches <u>NO</u> Water Marks <u>NO</u> Drift Lines <u>NO</u> Sediment Deposits <u>YES</u> Drainage Patterns in Wetlands</p> <p>Secondary Indicators</p> <p><u>YES</u> Oxidized Root Channels in Upper 12 Inches <u>NO</u> Water-Stained Leaves <u>NO</u> Local Soil Survey Data <u>YES</u> FAC-Neutral Test <u>NO</u> Other(Explain in Remarks)</p>
---	--

Remarks:

**DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)**

Project/Site: Big Spring Creek	Project No: Task 29	Date: 5-Aug-2004
Applicant/Owner: Montana Department of Transportation	County: Fergus	State: Montana
Investigators: Traxler	Plot ID: 1	

SOILS

Map Unit Name (Series and Phase): Enbar-Nesda loams, 0-2% slopes	Mapped Hydric Inclusion?
Map Symbol: 83 Drainage Class: SPD	Field Observations Confirm Mapped Type? <input checked="" type="radio"/> Yes <input type="radio"/> No
Taxonomy (Subgroup): Cumulic Haploborolls	

Depth (Inches)	Horizon	Matrix Color (Munsell Moist)	Mottle Color (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc
8	B	10YR3/1	N/A	N/A N/A	Silt loam
8	B	10YR3/2	10YR4/6	Common Distinct	Silt loam

Hydric Soil Indicators:

<u>NO</u> Histic Epipedon	<u>NO</u> Concretions
<u>NO</u> Sulfidic Odor	<u>NO</u> High Organic Content in Surface Layer in Sandy Soils
<u>NO</u> Aquic Moisture Regime	<u>NO</u> Organic Streaking in Sandy Soils
<u>NO</u> Reducing Conditions	<u>NO</u> Listed on Local Hydric Soils List
<u>YES</u> Gleyed or Low Chroma Colors	<u>NO</u> Listed on National Hydric Soils List
	<u>NO</u> Other (Explain in Remarks)

Remarks:

WETLAND DETERMINATION

Hydrophytic Vegetation Present? <input checked="" type="radio"/> Yes <input type="radio"/> No	Is the Sampling Point within the Wetland? <input checked="" type="radio"/> Yes <input type="radio"/> No
Wetland Hydrology Present? <input checked="" type="radio"/> Yes <input type="radio"/> No	
Hydric Soils Present? <input checked="" type="radio"/> Yes <input type="radio"/> No	

Remarks:
This plot was taken in apparent pre-existing wetland, east of the stream and in the south portion of the site.

DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

Project/Site: Big Spring Creek	Project No: Task 29	Date: 5-Aug-2004
Applicant/Owner: Montana Department of Transportation	County: Fergus	State: Montana
Investigators: Traxler	Plot ID: 2	

Project/Site: Big Spring Creek	Project No: Task 29	Date: 5-Aug-2004
Applicant/Owner: Montana Department of Transportation	County: Fergus	State: Montana
Investigators: Traxler	Plot ID: 2	

Do Normal Circumstances exist on the site? is the site significantly disturbed (Atypical Situation)? is the area a potential Problem Area? (If needed, explain on the reverse side)	<input checked="" type="radio"/> Yes <input type="radio"/> No <input type="radio"/> Yes <input checked="" type="radio"/> No	Community ID: EM Transect ID: NA Field Location: hwy slope toe, NE portion of site
--	--	---

Map Unit Name (Series and Phase): Fluvaquentic Haplaquolls, nearly level	Mapped Hydric Inclusion?
Map Symbol: 105 Drainage Class: PD	Field Observations Confirm Mapped Type? <input checked="" type="radio"/> Yes No
Taxonomy (Subgroup): Fluvaquentic Haplaquolls	
Profile Description	

VEGETATION (USFWS Region No. 9)					
Dominant Plant Species(Latin/Common)	Stratum	Indicator	Plant Species(Latin/Common)	Stratum	Indicator
<i>Typha latifolia</i>	Herb	OBL	<i>Polygonum lapathifolium</i>	Herb	FACW+
Cattail,Broad-Leaf			Willow-Weed		
<i>Agrostis alba</i>	Herb	FACW	<i>Juncus articulatus</i>	Herb	OBL
Redtop			Rush,Jointed		
<i>Allopecurus pratensis</i>	Herb	FACW	<i>Juncus ensifolius</i>	Herb	FACW
Foxtail,Meadow			Rush,Three-Stamen		
<i>Bidens cernua</i>	Herb	FACW+	<i>Echinochloa crusgalli</i>	Herb	FACW
Beggar-Ticks,Nodding			Grass,Barnyard		
<i>Epilobium ciliatum</i>	Herb	FACW-	<i>Glyceria elata</i>	Herb	FACW+
Willow-Herb,Hairy			Grass,Tall Manna		
<i>Eleocharis palustris</i>	Herb	OBL	<i>Juncus torreyi</i>	Herb	FACW
Spikerush,Creeping			Rush,Torrey's		
<i>Rumex crispus</i>	Herb	FACW	<i>Plantago major</i>	Herb	FAC+
Dock,Curly			Plantain,Common		
<i>Carex aquatilis</i>	Herb	OBL	<i>Salix exigua</i>	Shrub	OBL
Sedge,Water			Willow,Sandbar		

Depth (Inches)	Horizon	Matrix Color (Munsell Moist)	Mottle Color (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc
10	B	10YR3/1	N/A	N/A N/A	Silty clay loam

Hydric Soil Indicators:	<input type="checkbox"/> NO Histosol <input type="checkbox"/> NO Histic Epipedon <input type="checkbox"/> NO Sulfidic Odor <input type="checkbox"/> NO Aquic Moisture Regime <input type="checkbox"/> NO Reducing Conditions <input checked="" type="checkbox"/> YES Gleyed or Low Chroma Colors	<input type="checkbox"/> NO Concretions <input type="checkbox"/> NO High Organic Content in Surface Layer in Sandy Soils <input type="checkbox"/> NO Organic Streaking in Sandy Soils <input type="checkbox"/> NO Listed on Local Hydric Soils List <input type="checkbox"/> NO Reducing Conditions <input type="checkbox"/> NO Listed on National Hydric Soils List <input type="checkbox"/> NO Other (Explain in Remarks)
-------------------------	---	---

Percent of Dominant Species that are OBL, FACW or FAC: (excluding FAC-) 16/16 = 100.00%	FAC Neutral: 15/15 = 100.00%	Numeric Index: 28/16 = 1.75
Remarks: Salix scattered.		

Remarks:

Hydrophytic Vegetation Present? <input checked="" type="radio"/> Yes No	Is the Sampling Point within the Wetland? <input checked="" type="radio"/> Yes No
Wetland Hydrology Present? <input checked="" type="radio"/> Yes No	
Hydric Soils Present? <input checked="" type="radio"/> Yes No	

Remarks: Plot taken along toe of highway fill slope in NE corner of site. This area is developing rapidly into a functional wetland.

HYDROLOGY YES Recorded Data(Describe in Remarks): <input type="checkbox"/> NO Stream, Lake or Tide Gauge <input checked="" type="checkbox"/> YES Aerial Photographs <input type="checkbox"/> NO Other <input checked="" type="checkbox"/> NO No Recorded Data Field Observations Depth of Surface Water: = 2 (in.) Depth to Free Water in Pit: N/A (in.) Depth to Saturated Soil: N/A (in.)	Wetland Hydrology Indicators Primary Indicators <input checked="" type="checkbox"/> YES Inundated <input checked="" type="checkbox"/> YES Saturated in Upper 12 Inches <input type="checkbox"/> NO Water Marks <input type="checkbox"/> NO Drift Lines <input type="checkbox"/> NO Sediment Deposits <input checked="" type="checkbox"/> YES Drainage Patterns in Wetlands Secondary Indicators <input type="checkbox"/> NO Oxidized Root Channels in Upper 12 Inches <input type="checkbox"/> NO Water-Stained Leaves <input type="checkbox"/> NO Local Soil Survey Data <input checked="" type="checkbox"/> YES FAC-Neutral Test <input type="checkbox"/> NO Other(Explain in Remarks)
Remarks: Spring-fed; obvious groundwater connection.	

DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

Project/Site: Big Spring Creek	Project No: Task 29	Date: 5-Aug-2004
Applicant/Owner: Montana Department of Transportation		County: Fergus
Investigators: Traxler		State: Montana
		Plot ID: 3

Do Normal Circumstances exist on the site?	<input checked="" type="radio"/> Yes <input type="radio"/> No	Community ID: Transitional
Is the site significantly disturbed (Atypical Situation)?	<input type="radio"/> Yes <input checked="" type="radio"/> No	Transect ID: NA
Is the area a potential Problem Area? (If needed, explain on the reverse side)	<input type="radio"/> Yes <input checked="" type="radio"/> No	Field Location: W. of stream, S. of Parking lot @ bend

VEGETATION (USFWS Region No. 9)

Dominant Plant Species(Latin/Common)	Stratum	Indicator	Plant Species(Latin/Common)	Stratum	Indicator
<i>Cirsium arvense</i>	Herb	FACU+	<i>Sonchus arvensis</i>	Herb	FACU+
Thistle,Creeping			Sowthistle,Field		
<i>Equisetum arvense</i>	Herb	FAC	<i>Bromus inermis</i>	Herb	NI
Horsetail,Field			Brome, smooth		
<i>Phleum pratense</i>	Herb	FACU	<i>Alopecurus pratensis</i>	Herb	FACW
Timothy			Foxtail,Meadow		
<i>Trifolium fragiferum</i>	Herb	FACU	<i>Trifolium repens</i>	Herb	FACU+
Clover,Strawberry			Clover,White		
<i>Agrostis alba</i>	Herb	FACW			
Redtop					

Percent of Dominant Species that are OBL, FACW or FAC (excluding FAC-) 3/8 = 37.50%	FAC Neutral: 2/7 = 28.57%
	Numeric Index: 27/8 = 3.38

Remarks:
Does not satisfy hydrophytic vegetation criteria yet, but continues to transition in that direction.

HYDROLOGY

<u>NO</u> Recorded Data(Describe in Remarks): N/A Stream, Lake or Tide Gauge N/A Aerial Photographs N/A Other YES No Recorded Data	Wetland Hydrology Indicators Primary Indicators <u>NO</u> Inundated YES Saturated in Upper 12 Inches <u>NO</u> Water Marks <u>NO</u> Drift Lines <u>NO</u> Sediment Deposits <u>NO</u> Drainage Patterns in Wetlands Secondary Indicators <u>NO</u> Oxidized Root Channels in Upper 12 Inches <u>NO</u> Water-Stained Leaves <u>NO</u> Local Soil Survey Data <u>NO</u> FAC-Neutral Test <u>NO</u> Other(Explain in Remarks)
Field Observations Depth of Surface Water: N/A (in.) Depth to Free Water in Pit: N/A (in.) Depth to Saturated Soil: = 8 (in.)	

Remarks:
Saturated.

DATA FORM
ROUTINE WETLAND DETERMINATION
(1987 COE Wetlands Delineation Manual)

Project/Site: Big Spring Creek	Project No: Task 29	Date: 5-Aug-2004
Applicant/Owner: Montana Department of Transportation		County: Fergus
Investigators: Traxler		State: Montana
		Plot ID: 3

SOILS

Map Unit Name (Series and Phase): Enbar-Nesda loams, 0-2% slopes	Mapped Hydric Inclusion?
Map Symbol: 83 Drainage Class: SPD	Field Observations Confirm Mapped Type? <input checked="" type="radio"/> Yes <input type="radio"/> No
Taxonomy (Subgroup): Cumulic Haploborolls	

Depth (Inches)	Horizon	Matrix Color (Munsell Moist)	Mottle Color (Munsell Moist)	Mottle Abundance/Contrast	Texture, Concretions, Structure, etc
10	B	10YR3/2	10YR4/6	Few Faint	Silt loam

Hydric Soil Indicators:

<u>NO</u> Histosol	<u>NO</u> Concretions
<u>NO</u> Histic Epipedon	<u>NO</u> High Organic Content in Surface Layer in Sandy Soils
<u>NO</u> Sulfidic Odor	<u>NO</u> Organic Streaking in Sandy Soils
<u>NO</u> Aquic Moisture Regime	<u>NO</u> Listed on Local Hydric Soils List
<u>NO</u> Reducing Conditions	<u>NO</u> Listed on National Hydric Soils List
YES Gleyed or Low Chroma Colors	<u>NO</u> Other (Explain in Remarks)

Remarks:
Mottles faint: soils developing.

WETLAND DETERMINATION

Hydrophytic Vegetation Present? <input type="radio"/> Yes <input checked="" type="radio"/> No	Is the Sampling Point within the Wetland? <input type="radio"/> Yes <input checked="" type="radio"/> No
Wetland Hydrology Present? <input checked="" type="radio"/> Yes <input type="radio"/> No	
Hydric Soils Present? <input checked="" type="radio"/> Yes <input type="radio"/> No	

Remarks:
Plot take in inside bend of creek, south of parking lot. This area is currently technically upland, but will likely develop a prevalence of hydrophytic vegetation, based on hydrology and soils. Area is considered transitional.



14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS AND ANIMALS

i. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S _____
- Secondary habitat (list species) D S _____
- Incidental habitat (list species) D S Bald eagle.
- No usable habitat D S _____

ii. **Rating** (Based on the strongest habitat chosen in 14A(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point and Rating	---	---	---	---	---	.3 (L)	---

If documented, list the source (e.g., observations, records, etc.): _____

14B. HABITAT FOR PLANTS AND ANIMALS RATED AS S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM.

Do not include species listed in 14A(i).

i. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S _____
- Secondary habitat (list species) D S _____
- Incidental habitat (list species) D S N. leopard frog.
- No usable habitat D S _____

iii. **Rating** (Based on the strongest habitat chosen in 14B(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level:	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point and Rating	---	---	---	---	---	.1 (L)	---

If documented, list the source (e.g., observations, records, etc.): _____

14C. General Wildlife Habitat Rating

i. **Evidence of overall wildlife use in the AA:** (Check either substantial, moderate, or low)

Substantial (based on any of the following)

- observations of abundant wildlife #s or high species diversity (during any period)
- abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
- presence of extremely limiting habitat features not available in the surrounding area
- interviews with local biologists with knowledge of the AA

Low (based on any of the following)

- few or no wildlife observations during peak use periods
- little to no wildlife sign
- sparse adjacent upland food sources
- interviews with local biologists with knowledge of AA

Moderate (based on any of the following)

- observations of scattered wildlife groups or individuals or relatively few species during peak periods
- common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
- adequate adjacent upland food sources
- interviews with local biologists with knowledge of the AA

ii. **Wildlife Habitat Features** (Working from top to bottom, select appropriate AA attributes to determine the exceptional (E), high (H), moderate (M), or low (L) rating. Structural diversity is from #13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in terms of their percent composition in the AA (see #10). Duration of Surface Water: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; A= absent.

Structural Diversity (from #13)	<input type="checkbox"/> High								<input type="checkbox"/> Moderate								<input checked="" type="checkbox"/> Low			
Class Cover Distribution (all vegetated classes)	<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input checked="" type="checkbox"/> Even			
Duration of Surface Water in ≥ 10% of AA	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Low disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	H	--	--	--
High disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

iii. **Rating** (Using 14C(i) and 14C(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L) for this function.)

Evidence of Wildlife Use from 14C(i)	Wildlife Habitat Features Rating from 14C(ii)			
	<input type="checkbox"/> Exceptional	<input checked="" type="checkbox"/> High	<input type="checkbox"/> Moderate	<input type="checkbox"/> Low
Substantial	--	--	--	--
Moderate	--	.7 (M)	--	--
Low	--	--	--	--

Comments: _____

14D. GENERAL FISH/AQUATIC HABITAT RATING NA (proceed to 14E)

If the AA is not or was not historically used by fish due to lack of habitat, excessive gradient, then check the NA box above.

Assess if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [e.g. fish use is precluded by perched culvert or other barrier, etc.]. If fish use occurs in the AA but is not desired from a resource management perspective (e.g. fish use within an irrigation canal), then Habitat Quality [14D(i)] below should be marked as "Low", applied accordingly in 14D(ii) below, and noted in the comments.

i. **Habitat Quality** (Pick the appropriate AA attributes in matrix to pick the exceptional (E), high (H), moderate (M), or low (L) quality rating.)

Duration of Surface Water in AA	<input checked="" type="checkbox"/> Permanent/Perennial			<input type="checkbox"/> Seasonal / Intermittent			<input type="checkbox"/> Temporary / Ephemeral		
Cover - % of waterbody in AA containing cover objects (e.g. submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation)	>25%	10-25%	<10%	>25%	10-25%	<10%	>25%	10-25%	<10%
Shading - >75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities	--	--	--	--	--	--	--	--	--
Shading - 50 to 75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	--	--	--	--	--	--	--
Shading - < 50% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	M	--	--	--	--	--	--

ii. **Modified Habitat Quality:** Is fish use of the AA precluded or significantly reduced by a culvert, dike, other man-made structure or activity or is the waterbody included on the 'MDEQ list of waterbodies in need of TMDL development' with 'Probable Impaired Uses' listed as cold or warm water fishery or aquatic life support?

Y N If yes, reduce the rating from 14D(i) by one level and check the modified habitat quality rating: E H M L

iii. **Rating** (Use the conclusions from 14D(i) and 14D(ii) above and the matrix below to pick the functional point and rating of exceptional (E), high (H), moderate (M), or low (L).)

Types of Fish Known or Suspected Within AA	Modified Habitat Quality from 14D(ii)			
	<input type="checkbox"/> Exceptional	<input type="checkbox"/> High	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> Low
Native game fish	--	--	.7 (M)	--
Introduced game fish	--	--	--	--
Non-game fish	--	--	--	--
No fish	--	--	--	--

Comments: Sauger, Mnt. Whitefish present (MRIS 2002)

14E. FLOOD ATTENUATION NA (proceed to 14G)

Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA do not flooded from in-channel or overbank flow, check NA above.

i. **Rating** (Working from top to bottom, mark the appropriate attributes to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Estimated wetland area in AA subject to periodic flooding	<input type="checkbox"/> ≥ 10 acres			<input type="checkbox"/> <10, >2 acres			<input checked="" type="checkbox"/> ≤2 acres		
% of flooded wetland classified as forested, scrub/shrub, or both	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
AA contains no outlet or restricted outlet	--	--	--	--	--	--	--	--	.2 (L)
AA contains unrestricted outlet	--	--	--	--	--	--	--	--	--

ii. **Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA?** (check)

Y N Comments: Residences.

14F. SHORT AND LONG TERM SURFACE WATER STORAGE NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow. If no wetlands in the AA are subject to flooding or ponding, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Abbreviations: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral.

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding.	<input type="checkbox"/> >5 acre feet			<input type="checkbox"/> <5, >1 acre feet			<input checked="" type="checkbox"/> ≤1 acre foot		
Duration of surface water at wetlands within the AA	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Wetlands in AA flood or pond ≥ 5 out of 10 years	--	--	--	--	--	--	--	.3 (L)	--
Wetlands in AA flood or pond < 5 out of 10 years	--	--	--	--	--	--	--	--	--

Comments: _____

14G. SEDIMENT/NUTRIENT/TOXICANT RETENTION AND REMOVAL NA (proceed to 14H)

Applies to wetlands with potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Sediment, Nutrient, and Toxicant Input Levels Within AA	AA receives or surrounding land use has potential to deliver low to moderate levels of sediments, nutrients, or compounds such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
% cover of wetland vegetation in AA	<input type="checkbox"/> ≥ 70%		<input checked="" type="checkbox"/> < 70%		<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%	
Evidence of flooding or ponding in AA	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No
AA contains no or restricted outlet	--	--	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	.6 (M)	--	--	--	--	--

Comments: _____

14H. SEDIMENT/ShORELINE STABILIZATION

NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body that is subject to wave action. If this does not apply, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating exceptional (E), high (H), moderate (M), or low (L) for this function.

% Cover of wetland streambank or shoreline by species with deep, binding rootmasses.	Duration of Surface Water Adjacent to Rooted Vegetation		
	<input checked="" type="checkbox"/> Permanent / Perennial	<input type="checkbox"/> Seasonal / Intermittent	<input type="checkbox"/> Temporary / Ephemeral
≥ 65 %	--	--	--
35-64 %	.7 (M)	--	--
< 35 %	--	--	--

Comments: Planted shrubs will improve this rating once better established.

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

A = acreage of vegetated component in the AA. B = structural diversity rating from #13. C = Yes (Y) or No (N) as to whether or not the AA contains a surface or subsurface outlet; P/P = permanent/perennial; S/I = seasonal/intermittent; T/E/A = temporary/ephemeral/absent.

A	<input type="checkbox"/> Vegetated component >5 acres						<input type="checkbox"/> Vegetated component 1-5 acres						<input checked="" type="checkbox"/> Vegetated component <1 acre					
B	<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input checked="" type="checkbox"/> Low	
C	<input type="checkbox"/> Y	<input checked="" type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input checked="" type="checkbox"/> Y	<input type="checkbox"/> N
P/P	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	.4M	--
S/I	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
T/E/A	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Comments:

14J. GROUNDWATER DISCHARGE/RECHARGE (D/R) (Check the indicators in i & ii below that apply to the AA)

i. **Discharge Indicators**

- Springs are known or observed.
- Vegetation growing during dormant season/drought.
- Wetland occurs at the toe of a natural slopes.
- Seeps are present at the wetland edge.
- AA permanently flooded during drought periods.
- Wetland contains an outlet, but no inlet.
- Other Alluvial flow.

ii. **Recharge Indicators**

- Permeable substrate presents without underlying impeding layer.
- Wetland contains inlet but not outlet.
- Other _____

iii. **Rating:** Use the information from 14J(i) and 14J(ii) above and the table below to arrive at the functional point and rating of high (H) or low (L) for this function.

Criteria	Functional Point and Rating
AA has known Discharge/Recharge area or one or more indicators of D/R present	1 (H)
No Discharge/Recharge indicators present	--
Available Discharge/Recharge information inadequate to rate AA D/R potential	--

Comments: _____

14K. UNIQUENESS

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP.			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP.			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate.		
	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input checked="" type="checkbox"/> common	<input type="checkbox"/> abundant
Estimated Relative Abundance from #11									
Low disturbance at AA (#12i)	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (#12i)	--	--	--	--	--	--	--	.3L	--
High disturbance at AA (#12i)	--	--	--	--	--	--	--	--	--

Comments: _____

14L. RECREATION / EDUCATION POTENTIAL

i. Is the AA a known recreational or educational site? Yes (Rate High (1.0), then proceed to 14L(ii) only] No [Proceed to 14L(iii)]

ii. Check categories that apply to the AA: Educational / scientific study Consumptive rec. Non-consumptive rec. Other

iii. Based on the location, diversity, size, and other site attributes, is there a strong potential for recreational or educational use?

- Yes [Proceed to 14L (ii) and then 14L(iv).]
- No [Rate as low in 14L(iv)]

iv. **Rating** (Use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Ownership	Disturbance at AA from #12(i)	
	<input checked="" type="checkbox"/> Low	<input type="checkbox"/> High
Public ownership	1 (H)	--
Private ownership	--	--

Comments: Fishing, established Park, school nearby.

FUNCTION, VALUE SUMMARY, AND OVERALL RATING

Function and Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units (Actual Points x Estimated AA Acreage)
A. Listed/Proposed T&E Species Habitat	Low	0.30	1	
B. MT Natural Heritage Program Species Habitat	Low	0.10	1	
C. General Wildlife Habitat	Mod	0.70	1	
D. General Fish/Aquatic Habitat	Mod	0.70	1	
E. Flood Attenuation	Low	0.20	1	
F. Short and Long Term Surface Water Storage	Low	0.30	1	
G. Sediment/Nutrient/Toxicant Removal	Mod	0.60	1	
H. Sediment/Shoreline Stabilization	Mod	0.70	1	
I. Production Export/Food Chain Support	Mod	0.40	1	
J. Groundwater Discharge/Recharge	High	1.00	1	
K. Uniqueness	Low	0.30	1	
L. Recreation/Education Potential	High	1.00	1	
Totals:		<u>5.30</u>	<u>12.00</u>	
Percent of Total Possible Points:			44% (Actual / Possible) x 100 [rd to nearest whole #]	

Category I Wetland: (Must satisfy **one** of the following criteria. If not proceed to Category II.)

Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; **or**

Score of 1 functional point for Uniqueness; **or**

Score of 1 functional point for Flood Attenuation **and** answer to Question 14E(ii) is "yes"; **or**

Percent of total Possible Points is > 80%.

Category II Wetland: (Criteria for Category I not satisfied **and** meets any **one** of the following Category II criteria. If not satisfied, proceed to Category IV.)

Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; **or**

Score of .9 or 1 functional point for General Wildlife Habitat; **or**

Score of .9 or 1 functional point for General Fish/Aquatic Habitat; **or**

"High" to "Exceptional" ratings for **both** General Wildlife Habitat **and** General Fish / Aquatic Habitat; **or**

Score of .9 functional point for Uniqueness; **or**

Percent of total possible points is > 65%.

Category III Wetland: (Criteria for Categories I, II, or IV not satisfied.)

Category IV Wetland: (Criteria for Categories I or II are not satisfied **and** all of the following criteria are met; If not satisfied, proceed to Category III.)

"Low" rating for Uniqueness; **and**

"Low" rating for Production Export / Food Chain Support; **and**

Percent of total possible points is < 30%.

OVERALL ANALYSIS AREA (AA) RATING: (Check appropriate category based on the criteria outlined above.)

I
 II
 III
 IV

14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS AND ANIMALS

iv. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S _____
- Secondary habitat (list species) D S _____
- Incidental habitat (list species) D S _____
- No usable habitat D S _____

v. **Rating** (Based on the strongest habitat chosen in 14A(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point and Rating	---	---	---	---	---	---	0 (L)

If documented, list the source (e.g., observations, records, etc.): _____

14B. HABITAT FOR PLANTS AND ANIMALS RATED AS S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM.

Do not include species listed in 14A(i).

ii. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S _____
- Secondary habitat (list species) D S _____
- Incidental habitat (list species) D S N. leopard frog.
- No usable habitat D S _____

vi. **Rating** (Based on the strongest habitat chosen in 14B(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level:	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point and Rating	---	---	---	---	---	.1 (L)	---

If documented, list the source (e.g., observations, records, etc.): _____

14C. General Wildlife Habitat Rating

ii. **Evidence of overall wildlife use in the AA:** (Check either substantial, moderate, or low)

- Substantial** (based on any of the following)
 - observations of abundant wildlife #s or high species diversity (during any period)
 - abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
 - presence of extremely limiting habitat features not available in the surrounding area
 - interviews with local biologists with knowledge of the AA
- Low** (based on any of the following)
 - few or no wildlife observations during peak use periods
 - little to no wildlife sign
 - sparse adjacent upland food sources
 - interviews with local biologists with knowledge of AA
- Moderate** (based on any of the following)
 - observations of scattered wildlife groups or individuals or relatively few species during peak periods
 - common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
 - adequate adjacent upland food sources
 - interviews with local biologists with knowledge of the AA

ii. **Wildlife Habitat Features** (Working from top to bottom, select appropriate AA attributes to determine the exceptional (E), high (H), moderate (M), or low (L) rating. Structural diversity is from #13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in terms of their percent composition in the AA (see #10). Duration of Surface Water: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; A= absent.

Structural Diversity (from #13)	<input type="checkbox"/> High								<input type="checkbox"/> Moderate								<input checked="" type="checkbox"/> Low			
Class Cover Distribution (all vegetated classes)	<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input checked="" type="checkbox"/> Even			
Duration of Surface Water in ≥ 10% of AA	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Low disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	M	--	--
High disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

iii. **Rating** (Using 14C(i) and 14C(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L) for this function.)

Evidence of Wildlife Use from 14C(i)	Wildlife Habitat Features Rating from 14C(ii)			
	<input type="checkbox"/> Exceptional	<input type="checkbox"/> High	<input checked="" type="checkbox"/> Moderate	<input type="checkbox"/> Low
Substantial	--	--	--	--
Moderate	--	--	.5 (M)	--
ow	--	--	--	--

Comments: _____

14D. GENERAL FISH/AQUATIC HABITAT RATING NA (proceed to 14E)

If the AA is not or was not historically used by fish due to lack of habitat, excessive gradient, then check the NA box above.

Assess if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [e.g. fish use is precluded by perched culvert or other barrier, etc.]. If fish use occurs in the AA but is not desired from a resource management perspective (e.g. fish use within an irrigation canal), then Habitat Quality [14D(i)] below should be marked as "Low", applied accordingly in 14D(ii) below, and noted in the comments.

i. **Habitat Quality** (Pick the appropriate AA attributes in matrix to pick the exceptional (E), high (H), moderate (M), or low (L) quality rating.)

Duration of Surface Water in AA	<input type="checkbox"/> Permanent/Perennial			<input type="checkbox"/> Seasonal / Intermittent			<input type="checkbox"/> Temporary / Ephemeral		
Cover - % of waterbody in AA containing cover objects (e.g. submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation)	>25%	10-25%	<10%	>25%	10-25%	<10%	>25%	10-25%	<10%
Shading - >75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities	--	--	--	--	--	--	--	--	--
Shading - 50 to 75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	--	--	--	--	--	--	--
Shading - < 50% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	--	--	--	--	--	--	--

ii. **Modified Habitat Quality:** Is fish use of the AA precluded or significantly reduced by a culvert, dike, other man-made structure or activity or is the waterbody included on the 'MDEQ list of waterbodies in need of TMDL development' with 'Probable Impaired Uses' listed as cold or warm water fishery or aquatic life support? Y N If yes, reduce the rating from 14D(i) by one level and check the modified habitat quality rating: E H M L

iii. **Rating** (Use the conclusions from 14D(i) and 14D(ii) above and the matrix below to pick the functional point and rating of exceptional (E), high (H), moderate (M), or low (L).)

Types of Fish Known or Suspected Within AA	Modified Habitat Quality from 14D(ii)			
	<input type="checkbox"/> Exceptional	<input type="checkbox"/> High	<input type="checkbox"/> Moderate	<input type="checkbox"/> Low
Native game fish	--	--	--	--
Introduced game fish	--	--	--	--
Non-game fish	--	--	--	--
No fish	--	--	--	--

Comments: _____

14E. FLOOD ATTENUATION NA (proceed to 14G)

Applies only to wetlands subject to flooding via in-channel or overbank flow. If wetlands in AA do not flooded from in-channel or overbank flow, check NA above.

i. **Rating** (Working from top to bottom, mark the appropriate attributes to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Estimated wetland area in AA subject to periodic flooding	<input type="checkbox"/> ≥ 10 acres			<input type="checkbox"/> <10, >2 acres			<input checked="" type="checkbox"/> ≤2 acres		
% of flooded wetland classified as forested, scrub/shrub, or both	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
AA contains no outlet or restricted outlet	--	--	--	--	--	--	--	--	.2 (L)
AA contains unrestricted outlet	--	--	--	--	--	--	--	--	--

ii. **Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA?** (check) Y N Comments: Residences.

14F. SHORT AND LONG TERM SURFACE WATER STORAGE NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow. If no wetlands in the AA are subject to flooding or ponding, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.) Abbreviations: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral.

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding.	<input type="checkbox"/> >5 acre feet			<input type="checkbox"/> <5, >1 acre feet			<input checked="" type="checkbox"/> ≤1 acre foot		
Duration of surface water at wetlands within the AA	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Wetlands in AA flood or pond ≥ 5 out of 10 years	--	--	--	--	--	--	--	.3 (L)	--
Wetlands in AA flood or pond < 5 out of 10 years	--	--	--	--	--	--	--	--	--

Comments: _____

14G. SEDIMENT/NUTRIENT/TOXICANT RETENTION AND REMOVAL NA (proceed to 14H)

Applies to wetlands with potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input. If no wetlands in the AA are subject to such input, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Sediment, Nutrient, and Toxicant Input Levels Within AA	AA receives or surrounding land use has potential to deliver low to moderate levels of sediments, nutrients, or compounds such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
% cover of wetland vegetation in AA	<input checked="" type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%		<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%	
Evidence of flooding or ponding in AA	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No
AA contains no or restricted outlet	1 (H)	--	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	--	--	--	--	--	--

Comments: _____

14H. SEDIMENT/ShORELINE STABILIZATION

NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body that is subject to wave action. If this does not apply, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating exceptional (E), high (H), moderate (M), or low (L) for this function.

% Cover of wetland streambank or shoreline by species with deep, binding rootmasses.	Duration of Surface Water Adjacent to Rooted Vegetation		
	<input type="checkbox"/> Permanent / Perennial	<input type="checkbox"/> Seasonal / Intermittent	<input type="checkbox"/> Temporary / Ephemeral
≥ 65 %	--	--	--
35-64 %	--	--	--
< 35 %	--	--	--

Comments: _____

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

A = acreage of vegetated component in the AA. B = structural diversity rating from #13. C = Yes (Y) or No (N) as to whether or not the AA contains a surface or subsurface outlet; P/P = permanent/perennial; S/I = seasonal/intermittent; T/E/A = temporary/ephemeral/absent.

A	<input type="checkbox"/> Vegetated component >5 acres						<input type="checkbox"/> Vegetated component 1-5 acres						<input checked="" type="checkbox"/> Vegetated component <1 acre					
B	<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input checked="" type="checkbox"/> Low	
C	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input checked="" type="checkbox"/> Y	<input type="checkbox"/> N
P/P	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
S/I	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	.3L	--
T/E/A	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Comments: Subsurface.

14J. GROUNDWATER DISCHARGE/RECHARGE (D/R) (Check the indicators in i & ii below that apply to the AA)

i. **Discharge Indicators**

- Springs are known or observed.
- Vegetation growing during dormant season/drought.
- Wetland occurs at the toe of a natural slopes.
- Seeps are present at the wetland edge.
- AA permanently flooded during drought periods.
- Wetland contains an outlet, but no inlet.
- Other Alluvial flow.

ii. **Recharge Indicators**

- Permeable substrate presents without underlying impeding layer.
- Wetland contains inlet but not outlet.
- Other _____

iii. **Rating:** Use the information from 14J(i) and 14J(ii) above and the table below to arrive at the functional point and rating of high (H) or low (L) for this function.

Criteria	Functional Point and Rating
AA has known Discharge/Recharge area or one or more indicators of D/R present	1 (H)
No Discharge/Recharge indicators present	--
Available Discharge/Recharge information inadequate to rate AA D/R potential	--

Comments: _____

14K. UNIQUENESS

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP.			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP.			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate.		
	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input checked="" type="checkbox"/> common	<input type="checkbox"/> abundant
Estimated Relative Abundance from #11									
Low disturbance at AA (#12i)	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (#12i)	--	--	--	--	--	--	--	.3L	--
High disturbance at AA (#12i)	--	--	--	--	--	--	--	--	--

Comments: _____

14L. RECREATION / EDUCATION POTENTIAL

i. Is the AA a known recreational or educational site? Yes (Rate High (1.0), then proceed to 14L(ii) only] No [Proceed to 14L(iii)]

ii. Check categories that apply to the AA: Educational / scientific study Consumptive rec. Non-consumptive rec. Other

iii. Based on the location, diversity, size, and other site attributes, is there a strong potential for recreational or educational use?

- Yes [Proceed to 14L (ii) and then 14L(iv).]
- No [Rate as low in 14L(iv)]

iv. **Rating** (Use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Ownership	Disturbance at AA from #12(i)	
	<input type="checkbox"/> Low	<input checked="" type="checkbox"/> Moderate <input type="checkbox"/> High
Public ownership	--	.5(M)
Private ownership	--	--

Comments: School nearby, public site, moderate potential for study of wetland development.

FUNCTION, VALUE SUMMARY, AND OVERALL RATING

Function and Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units (Actual Points x Estimated AA Acreage)
A. Listed/Proposed T&E Species Habitat	Low	0.00	1	
B. MT Natural Heritage Program Species Habitat	Low	0.10	1	
C. General Wildlife Habitat	Mod	0.50	1	
D. General Fish/Aquatic Habitat	N/A	0.00	--	
E. Flood Attenuation	Low	0.20	1	
F. Short and Long Term Surface Water Storage	Low	0.30	1	
G. Sediment/Nutrient/Toxicant Removal	High	1.00	1	
H. Sediment/Shoreline Stabilization	N/A	0.00	--	
I. Production Export/Food Chain Support	Low	0.3	1	
J. Groundwater Discharge/Recharge	High	1.00	1	
K. Uniqueness	Low	0.30	1	
L. Recreation/Education Potential	Mod	0.50	1	
Totals:		<u>4.20</u>	<u>10.00</u>	
Percent of Total Possible Points:			42% (Actual / Possible) x 100 [rd to nearest whole #]	

<p>Category I Wetland: (Must satisfy one of the following criteria. If not proceed to Category II.)</p> <p><input type="checkbox"/> Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; or</p> <p><input type="checkbox"/> Score of 1 functional point for Uniqueness; or</p> <p><input type="checkbox"/> Score of 1 functional point for Flood Attenuation and answer to Question 14E(ii) is "yes"; or</p> <p><input type="checkbox"/> Percent of total Possible Points is > 80%.</p>
<p>Category II Wetland: (Criteria for Category I not satisfied and meets any one of the following Category II criteria. If not satisfied, proceed to Category IV.)</p> <p><input type="checkbox"/> Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; or</p> <p><input type="checkbox"/> Score of .9 or 1 functional point for General Wildlife Habitat; or</p> <p><input type="checkbox"/> Score of .9 or 1 functional point for General Fish/Aquatic Habitat; or</p> <p><input type="checkbox"/> "High" to "Exceptional" ratings for both General Wildlife Habitat and General Fish / Aquatic Habitat; or</p> <p><input type="checkbox"/> Score of .9 functional point for Uniqueness; or</p> <p><input type="checkbox"/> Percent of total possible points is > 65%.</p>
<p><input checked="" type="checkbox"/> Category III Wetland: (Criteria for Categories I, II, or IV not satisfied.)</p>
<p>Category IV Wetland: (Criteria for Categories I or II are not satisfied and <u>all</u> of the following criteria are met; If not satisfied, proceed to Category III.)</p> <p><input checked="" type="checkbox"/> "Low" rating for Uniqueness; and</p> <p><input checked="" type="checkbox"/> "Low" rating for Production Export / Food Chain Support; and</p> <p><input type="checkbox"/> Percent of total possible points is < 30%.</p>

OVERALL ANALYSIS AREA (AA) RATING: (Check appropriate category based on the criteria outlined above.)

I
 II
 III
 IV

14A. HABITAT FOR FEDERALLY LISTED OR PROPOSED THREATENED OR ENDANGERED PLANTS AND ANIMALS

vii. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S
- Secondary habitat (list species) D S
- Incidental habitat (list species) D S bald eagle
- No usable habitat D S

viii. **Rating** (Based on the strongest habitat chosen in 14A(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point and Rating	---	---	---	---	---	.3 (L)	---

If documented, list the source (e.g., observations, records, etc.): _____

14B. HABITAT FOR PLANTS AND ANIMALS RATED AS S1, S2, OR S3 BY THE MONTANA NATURAL HERITAGE PROGRAM.

Do not include species listed in 14A(i).

iii. AA is Documented (D) or Suspected (S) to contain (check box):

- Primary or Critical habitat (list species) D S _____
- Secondary habitat (list species) D S N. leopard frog, possible ws cutthroat
- Incidental habitat (list species) D S _____
- No usable habitat D S _____

ix. **Rating** (Based on the strongest habitat chosen in 14B(i) above, find the corresponding rating of High (H), Moderate (M), or Low (L) for this function.

Highest Habitat Level:	doc/primary	sus/primary	doc/secondary	sus/secondary	doc/incidental	sus/incidental	none
Functional Point and Rating	---	---	---	.6 (M)	---	---	---

If documented, list the source (e.g., observations, records, etc.): _____

14C. General Wildlife Habitat Rating

iii. **Evidence of overall wildlife use in the AA:** (Check either substantial, moderate, or low)

- Substantial** (based on any of the following)
 - observations of abundant wildlife #s or high species diversity (during any period)
 - abundant wildlife sign such as scat, tracks, nest structures, game trails, etc.
 - presence of extremely limiting habitat features not available in the surrounding area
 - interviews with local biologists with knowledge of the AA
- Low** (based on any of the following)
 - few or no wildlife observations during peak use periods
 - little to no wildlife sign
 - sparse adjacent upland food sources
 - interviews with local biologists with knowledge of AA
- Moderate** (based on any of the following)
 - observations of scattered wildlife groups or individuals or relatively few species during peak periods
 - common occurrence of wildlife sign such as scat, tracks, nest structures, game trails, etc.
 - adequate adjacent upland food sources
 - interviews with local biologists with knowledge of the AA

ii. **Wildlife Habitat Features** (Working from top to bottom, select appropriate AA attributes to determine the exceptional (E), high (H), moderate (M), or low (L) rating. Structural diversity is from #13. For class cover to be considered evenly distributed, vegetated classes must be within 20% of each other in terms of their percent composition in the AA (see #10). Duration of Surface Water: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral; A= absent.

Structural Diversity (from #13)	<input type="checkbox"/> High								<input checked="" type="checkbox"/> Moderate								<input type="checkbox"/> Low			
Class Cover Distribution (all vegetated classes)	<input type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input checked="" type="checkbox"/> Even				<input type="checkbox"/> Uneven				<input type="checkbox"/> Even			
Duration of Surface Water in ≥ 10% of AA	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A	P/P	S/I	T/E	A
Low disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (see #12)	--	--	--	--	--	--	--	--	H	--	--	--	--	--	--	--	--	--	--	--
High disturbance at AA (see #12)	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

iii. **Rating** (Using 14C(i) and 14C(ii) above and the matrix below to arrive at the functional point and rating of exceptional (E), high (H), moderate (M), or low (L) for this function.)

Evidence of Wildlife Use from 14C(i)	Wildlife Habitat Features Rating from 14C(ii)			
	<input type="checkbox"/> Exceptional	<input checked="" type="checkbox"/> High	<input type="checkbox"/> Moderate	<input type="checkbox"/> Low
Substantial	--	.9 (H)	--	--
Moderate	--	--	--	--
Low	--	--	--	--

Comments: _____

14D. GENERAL FISH/AQUATIC HABITAT RATING NA (proceed to 14E)

If the AA is not or was not historically used by fish due to lack of habitat, excessive gradient, then check the NA box above.

Assess if the AA is used by fish or the existing situation is "correctable" such that the AA could be used by fish [e.g. fish use is precluded by perched culvert or other barrier, etc.]. If fish use occurs in the AA but is not desired from a resource management perspective (e.g. fish use within an irrigation canal), then Habitat Quality [14D(i)] below should be marked as "Low", applied accordingly in 14D(ii) below, and noted in the comments.

i. **Habitat Quality** (Pick the appropriate AA attributes in matrix to pick the exceptional (E), high (H), moderate (M), or low (L) quality rating.)

Duration of Surface Water in AA	<input checked="" type="checkbox"/> Permanent/Perennial			<input type="checkbox"/> Seasonal / Intermittent			<input type="checkbox"/> Temporary / Ephemeral		
Cover - % of waterbody in AA containing cover objects (e.g. submerged logs, large rocks & boulders, overhanging banks, floating-leaved vegetation)	>25%	10-25%	<10%	>25%	10-25%	<10%	>25%	10-25%	<10%
Shading - >75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities	--	--	--	--	--	--	--	--	--
Shading - 50 to 75% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	H	--	--	--	--	--	--	--
Shading - < 50% of streambank or shoreline of AA contains riparian or wetland scrub-shrub or forested communities.	--	--	--	--	--	--	--	--	--

ii. **Modified Habitat Quality:** Is fish use of the AA precluded or significantly reduced by a culvert, dike, other man-made structure or activity or is the waterbody included on the 'MDEQ list of waterbodies in need of TMDL development' with 'Probable Impaired Uses' listed as cold or warm water fishery or aquatic life support?
 Y N If yes, reduce the rating from 14D(i) by one level and check the modified habitat quality rating: E H M L

iii. **Rating** (Use the conclusions from 14D(i) and 14D(ii) above and the matrix below to pick the functional point and rating of exceptional (E), high (H), moderate (M), or low (L).)

Types of Fish Known or Suspected Within AA	Modified Habitat Quality from 14D(ii)			
	<input type="checkbox"/> Exceptional	<input checked="" type="checkbox"/> High	<input type="checkbox"/> Moderate	<input type="checkbox"/> Low
Native game fish	--	.9 (H)	--	--
Introduced game fish	--	--	--	--
Non-game fish	--	--	--	--
No fish	--	--	--	--

Comments: Mtn. Whitefish, Sauger present (MRIS 2002)

14E. FLOOD ATTENUATION NA (proceed to 14G)

Applies only to wetlands subject to flooding via in-channel or overbank flow.
 If wetlands in AA do not flooded from in-channel or overbank flow, check NA above.

i. **Rating** (Working from top to bottom, mark the appropriate attributes to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Estimated wetland area in AA subject to periodic flooding	<input type="checkbox"/> ≥ 10 acres			<input checked="" type="checkbox"/> <10, >2 acres			<input type="checkbox"/> ≤2 acres		
% of flooded wetland classified as forested, scrub/shrub, or both	75%	25-75%	<25%	75%	25-75%	<25%	75%	25-75%	<25%
AA contains no outlet or restricted outlet	--	--	--	--	.7 (H)	--	--	--	--
AA contains unrestricted outlet	--	--	--	--	--	--	--	--	--

ii. **Are residences, businesses, or other features which may be significantly damaged by floods located within 0.5 miles downstream of the AA?** (check)
 Y N Comments: Residences

14F. SHORT AND LONG TERM SURFACE WATER STORAGE NA (proceed to 14G)

Applies to wetlands that flood or pond from overbank or in-channel flow, precipitation, upland surface flow, or groundwater flow.
 If no wetlands in the AA are subject to flooding or ponding, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)
 Abbreviations: P/P = permanent/perennial; S/I = seasonal/intermittent; T/E = temporary/ephemeral.

Estimated maximum acre feet of water contained in wetlands within the AA that are subject to periodic flooding or ponding.	<input type="checkbox"/> >5 acre feet			<input checked="" type="checkbox"/> <5, >1 acre feet			<input type="checkbox"/> ≤1 acre foot		
Duration of surface water at wetlands within the AA	P/P	S/I	T/E	P/P	S/I	T/E	P/P	S/I	T/E
Wetlands in AA flood or pond ≥ 5 out of 10 years	--	--	--	--	.6 (M)	--	--	--	--
Wetlands in AA flood or pond < 5 out of 10 years	--	--	--	--	--	--	--	--	--

Comments: _____

14G. SEDIMENT/NUTRIENT/TOXICANT RETENTION AND REMOVAL NA (proceed to 14H)

Applies to wetlands with potential to receive excess sediments, nutrients, or toxicants through influx of surface or ground water or direct input.
 If no wetlands in the AA are subject to such input, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.)

Sediment, Nutrient, and Toxicant Input Levels Within AA	AA receives or surrounding land use has potential to deliver low to moderate levels of sediments, nutrients, or compounds such that other functions are not substantially impaired. Minor sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.				Waterbody on MDEQ list of waterbodies in need of TMDL development for "probable causes" related to sediment, nutrients, or toxicants or AA receives or surrounding land use has potential to deliver high levels of sediments, nutrients, or compounds such that other functions are substantially impaired. Major sedimentation, sources of nutrients or toxicants, or signs of eutrophication present.			
% cover of wetland vegetation in AA	<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%		<input type="checkbox"/> ≥ 70%		<input type="checkbox"/> < 70%	
Evidence of flooding or ponding in AA	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No	<input type="checkbox"/> Yes	<input type="checkbox"/> No
AA contains no or restricted outlet	1 (H)	--	--	--	--	--	--	--
AA contains unrestricted outlet	--	--	--	--	--	--	--	--

Comments: _____

14H. SEDIMENT/ShORELINE STABILIZATION

NA (proceed to 14I)

Applies only if AA occurs on or within the banks of a river, stream, or other natural or man-made drainage, or on the shoreline of a standing water body that is subject to wave action. If this does not apply, check NA above.

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating exceptional (E), high (H), moderate (M), or low (L) for this function.

% Cover of wetland streambank or shoreline by species with deep, binding rootmasses.	Duration of Surface Water Adjacent to Rooted Vegetation		
	<input checked="" type="checkbox"/> Permanent / Perennial	<input type="checkbox"/> Seasonal / Intermittent	<input type="checkbox"/> Temporary / Ephemeral
≥ 65 %	--	--	--
35-64 %	.7 (M)	--	--
< 35 %	--	--	--

Comments: May increase with plantings.

14I. PRODUCTION EXPORT / FOOD CHAIN SUPPORT

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

A = acreage of vegetated component in the AA. B = structural diversity rating from #13. C = Yes (Y) or No (N) as to whether or not the AA contains a surface or subsurface outlet; P/P = permanent/perennial; S/I = seasonal/intermittent; T/E/A = temporary/ephemeral/absent.

A	<input checked="" type="checkbox"/> Vegetated component >5 acres						<input type="checkbox"/> Vegetated component 1-5 acres						<input type="checkbox"/> Vegetated component <1 acre					
B	<input type="checkbox"/> High		<input checked="" type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low		<input type="checkbox"/> High		<input type="checkbox"/> Moderate		<input type="checkbox"/> Low	
C	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input checked="" type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N	<input type="checkbox"/> Y	<input type="checkbox"/> N
P/P	--	--	--	.9H	--	--	--	--	--	--	--	--	--	--	--	--	--	--
S/I	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
T/E/A	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Comments:

14J. GROUNDWATER DISCHARGE/RECHARGE (D/R) (Check the indicators in i & ii below that apply to the AA)

i. **Discharge Indicators**

- Springs are known or observed.
- Vegetation growing during dormant season/drought.
- Wetland occurs at the toe of a natural slopes.
- Seeps are present at the wetland edge.
- AA permanently flooded during drought periods.
- Wetland contains an outlet, but no inlet.
- Other

ii. **Recharge Indicators**

- Permeable substrate presents without underlying impeding layer.
- Wetland contains inlet but not outlet.
- Other

iii. **Rating:** Use the information from 14J(i) and 14J(ii) above and the table below to arrive at the functional point and rating of high (H) or low (L) for this function.

Criteria	Functional Point and Rating
AA has known Discharge/Recharge area or one or more indicators of D/R present	1 (H)
No Discharge/Recharge indicators present	--
Available Discharge/Recharge information inadequate to rate AA D/R potential	--

Comments:

14K. UNIQUENESS

i. **Rating** (Working from top to bottom, use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Replacement Potential	AA contains fen, bog, warm springs or mature (>80 yr-old) forested wetland or plant association listed as "S1" by the MTNHP.			AA does not contain previously cited rare types and structural diversity (#13) is high or contains plant association listed as "S2" by the MTNHP.			AA does not contain previously cited rare types or associations and structural diversity (#13) is low-moderate.		
	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input type="checkbox"/> common	<input type="checkbox"/> abundant	<input type="checkbox"/> rare	<input checked="" type="checkbox"/> common	<input type="checkbox"/> abundant
Estimated Relative Abundance from #11	--	--	--	--	--	--	--	--	--
Low disturbance at AA (#12i)	--	--	--	--	--	--	--	--	--
Moderate disturbance at AA (#12i)	--	--	--	--	--	--	--	.3L	--
High disturbance at AA (#12i)	--	--	--	--	--	--	--	--	--

Comments:

14L. RECREATION / EDUCATION POTENTIAL

i. Is the AA a known recreational or educational site? Yes (Rate High (1.0), then proceed to 14L(ii) only] No [Proceed to 14L(iii)]

ii. Check categories that apply to the AA: Educational / scientific study Consumptive rec. Non-consumptive rec. Other

iii. Based on the location, diversity, size, and other site attributes, is there a strong potential for recreational or educational use?

- Yes [Proceed to 14L (ii) and then 14L(iv).]
- No [Rate as low in 14L(iv)]

iv. **Rating** (Use the matrix below to arrive at the functional point and rating of high (H), moderate (M), or low (L) for this function.

Ownership	Disturbance at AA from #12(i)	
	<input checked="" type="checkbox"/> Low	<input type="checkbox"/> High
Public ownership	1 (H)	--
Private ownership	--	--

Comments: _____

FUNCTION, VALUE SUMMARY, AND OVERALL RATING

Function and Value Variables	Rating	Actual Functional Points	Possible Functional Points	Functional Units (Actual Points x Estimated AA Acreage)
A. Listed/Proposed T&E Species Habitat	Low	0.30	1	
B. MT Natural Heritage Program Species Habitat	Mod	0.60	1	
C. General Wildlife Habitat	High	0.90	1	
D. General Fish/Aquatic Habitat	High	0.90	1	
E. Flood Attenuation	High	0.70	1	
F. Short and Long Term Surface Water Storage	Mod	0.60	1	
G. Sediment/Nutrient/Toxicant Removal	High	1.00	1	
H. Sediment/Shoreline Stabilization	Mod	0.70	1	
I. Production Export/Food Chain Support	High	0.90	1	
J. Groundwater Discharge/Recharge	High	1.00	1	
K. Uniqueness	Low	0.30	1	
L. Recreation/Education Potential	High	1.00	1	
Totals:		8.90	12.00	
Percent of Total Possible Points:			74% (Actual / Possible) x 100 [rd to nearest whole #]	

Category I Wetland: (Must satisfy **one** of the following criteria. If not proceed to Category II.)

Score of 1 functional point for Listed/Proposed Threatened or Endangered Species; **or**

Score of 1 functional point for Uniqueness; **or**

Score of 1 functional point for Flood Attenuation **and** answer to Question 14E(ii) is "yes"; **or**

Percent of total Possible Points is > 80%.

Category II Wetland: (Criteria for Category I not satisfied **and** meets any **one** of the following Category II criteria. If not satisfied, proceed to Category IV.)

Score of 1 functional point for Species Rated S1, S2, or S3 by the MT Natural Heritage Program; **or**

Score of .9 or 1 functional point for General Wildlife Habitat; **or**

Score of .9 or 1 functional point for General Fish/Aquatic Habitat; **or**

"High" to "Exceptional" ratings for **both** General Wildlife Habitat **and** General Fish / Aquatic Habitat; **or**

Score of .9 functional point for Uniqueness; **or**

Percent of total possible points is > 65%.

Category III Wetland: (Criteria for Categories I, II, or IV not satisfied.)

Category IV Wetland: (Criteria for Categories I or II are not satisfied **and** all of the following criteria are met; If not satisfied, proceed to Category III.)

"Low" rating for Uniqueness; **and**

"Low" rating for Production Export / Food Chain Support; **and**

Percent of total possible points is < 30%.

OVERALL ANALYSIS AREA (AA) RATING: (Check appropriate category based on the criteria outlined above.)

I II III IV

Appendix C

REPRESENTATIVE PHOTOGRAPHS 2004 AERIAL PHOTOGRAPH

*MDT Wetland Mitigation Monitoring
Big Spring Creek
Lewistown, Montana*



Photo Point 1: 346 degrees North
New Big Spring Creek channel



Photo Point 1: 300 degrees NW



Photo Point 1: 260 degrees West
New Big Spring Creek channel



Photo Point 2: 155 degrees SE
Location of old creek channel parallel to highway



Photo Point 3: 190 degrees SW



Photo Point 3: 340 Degrees North



Photo Point 4: 15 degrees NE
From center of walkway – 6 feet from west bridge end

Photo Point 4: 200 degrees SW
From center of walkway – 6 feet from west bridge end



Photo Point 5: 10 Degrees North
Photo looking North towards foot bridge

Photo Point 5: 100 degrees East



Vegetation Transect start: 94 degrees East

Vegetation Transect End: 274 degrees West

Big Spring Creek 2004 Aerial Photograph




Appendix D

CONCEPTUAL SITE LAYOUT

*MDT Wetland Mitigation Monitoring
Big Spring Creek
Lewistown, Montana*



Scale (feet)



LEGEND

-  Existing Channel
-  Existing Wetlands
-  Existing Open Water
-  Water Pipe Lines
-  Fence Line
-  Highway Right-of-Way
-  Overhead Telephone Lines
-  Proposed Channel with Center Line
-  Levels of Disturbance
-  Mature Vegetation to Remain as Riparian Wetland
-  Open Water Wetland to Remain
-  Wetland to be Created
-  Emergent Wetland to Remain

Lowest point approx. 1 mile

DESIGNED BY J. R. ...	DATE 01/01/88	PROJECT Final Design for the Restoration of the Spring Creek and the Associated Flowage at Browns Lake
CHECKED BY ...	DATE 01/01/88	
DRAWN BY ...	DATE 01/01/88	
INTER-FLUVE, INC.		
WETLAND PLAN		
INTER-FLUVE, INC.		
		54

Appendix E

BIRD SURVEY PROTOCOL GPS PROTOCOL

*MDT Wetland Mitigation Monitoring
Big Spring Creek
Lewistown, Montana*

BIRD SURVEY PROTOCOL

The following is an outline of the MDT Wetland Mitigation Site Monitoring Bird Survey Protocol. Though each site is vastly different, the bird survey data collection methods must be standardized to a certain degree to increase repeatability. An Area Search within a restricted time frame will be used to collect the following data: a bird species list, density, behavior, and habitat-type use. There will be some decisions that team members must make to fit the protocol to their particular site. Each of the following sections and the desired result describes the protocol established to reflect bird species use over time.

Species Use within the Mitigation Wetland: Survey Method

Result: To conduct a bird survey of the wetland mitigation site within a restricted period of time and the budget allotment.

Sites that can be circumambulated or walked throughout.

These types of sites will include ponds, enhanced historic river channels, wet meadows, and any area that can be surveyed from the entirety of its perimeter or walked throughout. If the wetland is not uncomfortably inundated, conduct several “meandering” transects through the site in an orderly fashion (record the number and approximate location/direction of the transects in the field notebook; they do not have to be formalized or staked). If a very small portion of the site cannot be crossed due to inundation, this method will also apply. Though the sizes of the site vary, each site will require surveying to the fullest extent possible within a set time limit. The optimum times to conduct the survey are in the morning hours. Conduct the survey from sunrise to no later than 11:00 AM. (Note: some sites may have to be surveyed in the late afternoon or evening due to time constraints or weather; if this is the case, record the time of day and include this information in your report discussion.) If the survey is completed before 11:00 AM and no additions are being made to the list, then the task is complete. The overall limiting factor regarding the number of hours that are spent conducting this survey is the number of budgeted hours; this determination must be made by site by each individual.

In many cases, binoculars will be the only instrument that is needed to identify and count the birds using the wetland. If the wetland includes deep water habitat that can not be assessed with binoculars, then a scope and tripod are necessary. If this is the case, establish as many lookout posts as necessary from key vantage points to collect the data. Depending on the size of the open water, more time may be spent viewing the mitigation area from these vantage points than is spent walking the peripheries of more shallow-water wetlands.

Sites that cannot be circumambulated.

These types of sites will include large-bodied waters, such as reservoirs, particularly those with deep water habitat (>6 ft) close to the shore and no wetland development in that area of the shoreline. If one area of the reservoir was graded in such a way to create or enhance the development of a wetland, then that will be the area in which the ambulatory bird survey is

conducted. The team member must then determine the length of the shoreline that will be surveyed during each visit.

As stated above in the ambulatory site section, these large sites most likely will have to be surveyed from established vantage points.

Species Use within the Mitigation Wetland: Data Recording

Result: A complete list of bird species using the site, an estimate of bird densities and associated behaviors, and identification of habitat use.

1. Bird Species List

Record the bird species on the Bird Survey - Field Data Sheet using the appropriate 4-letter code of the common name. The coding uses the first two letters of the first two words of the birds' common name or if one name, the first four (4) letters. For example, mourning dove is coded MODO and mallard is MALL. If an unknown individual is observed, use the following protocol and define your abbreviation at the bottom of the field data sheet: unknown shorebird: UNSB; unknown brown bird (UNBR); unknown warbler (UNWA); unknown waterfowl (UNWF). For a flyover of a flock of unknown species, use a term that describes the birds' general characteristics and include the approximate flock size in parentheses; do not fill in the habitat column. For example, a flock of black, medium-sized birds could be coded: UNBB / FO (25). You may also note on the data sheet if that particular individual is using a constructed nest box.

2. Bird Density

In the office, sum the Bird Survey – Field Data Sheet data by species and by behavior. Record this data in the Bird Summary Table.

3. Bird Behavior

Bird behavior must be identified by what is known. When a species is simply observed, the behavior that it is immediately exhibiting is what is recorded. Only behaviors that have discreet descriptive terms should be used. The following terms are recommended: breeding pair individual (BP); foraging (F); flyover (FO); loafing (L; e.g. sleeping, roosting, floating with head tucked under wing are loafing behaviors); and, nesting (N). If more behaviors are observed that do have a specific descriptive word, use them and we will add it to the protocol; descriptive words or phrases such as “migrating” or “living on site” are unknown behaviors.

4. Bird Species Habitat Use

We are interested in what bird species are using which particular habitat within the mitigation wetlands. This data is easily collected by simply recording what habitat the species was initially observed. Use the following broad category habitat classifications: aquatic bed (AB - rooted floating, floating-leaved, or submergent vegetation); forested (FO); marsh (MA – cattail, bulrush, emergent vegetation, etc. with surface water); open water (OW – primarily unvegetated); scrub-shrub (SS); and upland buffer (UP); wet meadow (WM – sedges, rushes, grasses with little to no surface water). If other categories are observed onsite that are not suggested here, we will make a new category next year.

GPS Mapping and Aerial Photo Referencing Procedure

The wetland boundaries, photograph location points and sampling locations were field located with mapping grade Trimble Geo III GPS units. The data was collected with a minimum of three positions per feature using Course/Acquisition code. The collected data was then transferred to a PC and differentially corrected to the nearest operating Community Base Station. The corrected data was then exported to ACAD drawings in Montana State Plain Coordinates NAD 83 international feet.

The GPS positions collected and processed had a 68% accuracy of 7 feet except in isolated areas of Tasks .008 and .011, where it went to 12 feet. This is within the 1 to 5 meter range listed as the expected accuracy of the mapping grade Trimble GPS.

Aerial reference points were used to position the aerial photographs. This positioning did not remove the distortion inherent in all photos; this imagery is to be used as a visual aide only. The located wetland boundaries were given a final review by the wetland biologist and adjustments were made if necessary.

Any relationship of features located to easement or property lines are not to be construed from these figures. These relationships can only be determined with a survey by a licensed surveyor.

Appendix F

MACROINVERTEBRATE SAMPLING PROTOCOL AND DATA

*MDT Wetland Mitigation Monitoring
Big Spring Creek
Lewistown, Montana*

AQUATIC INVERTEBRATE SAMPLING PROTOCOL

Equipment List

- D-frame sampling net with 1 mm mesh. Wildco is a good source of these.
- Spare net.
- 1-liter plastic sample jars, wide-mouth. VWR has these: catalog #36319-707.
- 95% ethanol: Northwest Scientific in Billings carries this.

All these other things are generally available at hardware or sporting goods stores. Make the labels on an ink jet printer preferably.

- hip waders.
- pre-printed sample labels (printed on Rite-in-the-Rain or other coated paper, two labels per sample).
- pencil.
- plastic pail (3 or 5 gallon).
- large tea strainer or framed screen.
- towel.
- tape for affixing label to jar.
- cooler with ice for sample storage.

Site Selection

Select the sampling site with these considerations in mind:

- Select a site accessible with hip waders. If substrates are too soft, lay a wide board down to walk on.
- Determine a location that is representative of the overall condition of the wetland.

Sampling

Wetland invertebrates inhabit the substrate, the water column, the stems and leaves of aquatic vegetation, and the water surface. Your goal is to sweep the collecting net through each of these habitat types, and then to combine the resulting samples into the 1-liter sample jar.

Dip out about a gallon of water into the pail. Pour about a cup of ethanol into the sample jar. Fill out the top half of the sample labels, using pencil, since ink will dissolve in the ethanol.

Ideally, you can sample a swath of water column from near-shore outward to a depth of approximately 3 feet with a long sweep of the net, keeping the net at about half the depth of the water throughout the sweep. Sweep the water surface as well. Pull the net through a vegetated area, beneath the water surface, for at least a meter of distance.

Sample the substrate by pulling the net along the bottom, bumping it against the substrate several times as you pull.



This step is optional, but it gives you a chance to see that you've collected some invertebrates. Rinse the net into the bucket, and look for insects, crustaceans, etc. If necessary, repeat the sampling process in a nearby location, and add the net contents to the bucket. Remember to sample all four environments.

Sieve the contents of the bucket through the straining device and pour or carefully scrape the contents of the strainer into the sample jar.

If you skip the bucket-and-sieve steps, simply lift handfuls of material out of the sampling net into the jars. In either case, please include some muck or mud and some vegetation in the jar. Often, you will have collected a large amount of vegetable material. If this is the case, lift out handfuls of material from the sieve into the jar, until the jar is about half full. Please limit material you include in the sample, so that there is only a single jar for each sample.

Top off the sample jar with enough ethanol to cover all the material in the jar. Leave as little headroom as possible.

It is not necessary to sample habitats in any specified order. Keep in mind that disturbing the habitats prior to sampling will chase off the animals you are trying to capture.

Complete the sample labels. Place one label inside the sample jar and tape the other label securely to the outside of the jar. Dry the jar before attaching the outer label if necessary. In some situations, it may be necessary to collect more than one sample at a site. If you take multiple samples from the same site, clearly indicate this by using individual sample numbers, along with the total number of samples collected at the site (e.g. Sample #3 of 5 total samples).

Photograph the sampled site.

Sample Handling/Shipping

- In the field, keep collected samples cool by storing them in a cooler. Only a small amount of ice is necessary.
- Inventory all samples, preparing a list of all sites and enumerating all samples, before shipping or delivering to the laboratory.
- Deliver samples to Rhithron.

**MDT Wetland Mitigation Monitoring Project
Aquatic Invertebrate Monitoring
Summary 2001 - 2004**

METHODS

Among other monitoring activities, aquatic invertebrate assemblages were collected at a number of mitigation wetlands throughout Montana. This report summarizes data generated from four years of collection.

The method employed to assess these wetlands is based on constructing an index using a battery of 12 bioassessment metrics or attributes (Table 1) tested and recommended by Stribling et al. (1995) in a report to the Montana Department of Health and Environmental Science. In that study, it was determined that some of the metrics were of limited use in some geographic regions, and for some wetland types. Despite that finding, all 12 metrics are used in this evaluation of mitigated wetlands, since detailed geographic information and wetland classifications were unavailable.

Scoring criteria for metrics were developed by generally following the tactic used by Stribling et al. Boxplots were generated using a statistical software package, and distributions, median values, ranges, and quartiles for each metric were examined. All sites in all years of sampling were used. Camp Creek, which was sampled in 2002, 2003, and 2004, was assessed using the tested metric battery developed for montane streams of Western Montana (Bollman 1998). The fauna at the Camp Creek site was different from that of the other sites, and suggested montane stream conditions rather than wetland conditions. For the wetlands, "optimal" scores were generally those that fell above the 75th percentile (for those metrics that decrease in value in response to stress) or below the 25th percentile (for metrics that respond to stress by an increase in value) of all scores. Additional scoring ranges were established by bisecting the range below the 75th percentile for decreasing scores (or above the 25th percentile for increasing scores) into "sub-optimal" and "poor" assessment categories. A score of 5, 3, or 1 was assigned to optimal, sub-optimal, and poor metric performance, respectively. In this way, metric values were translated into normalized metric scores, and scores for all metrics were summed to produce a total bioassessment score. Total bioassessment scores were classified according to a similar process, using the ranges and distributions of total scores for all sites studied in all years.

The purpose of constructing an index from biological attributes or metrics is to provide a means of integrating information to facilitate the determination of whether management action is needed. The nature of the action needed is not determined solely by the index score, however, but by consideration of an analysis of the component metrics, the taxonomic composition of the assemblages, and other issues. The diagnostic functions of the metrics and taxonomic data need more study; our understanding of the interrelationships of natural environmental factors and anthropogenic disturbances are tentative. Thus, the further interpretive remarks accompanying the raw taxonomic and metric data are offered cautiously.

Sample processing

Aquatic invertebrate samples were collected at mitigation wetland sites in the summer months of 2001, 2002, 2003, and 2004 by personnel of Land and Water Consulting, Inc. Sampling procedures utilized were based on the protocols developed by the Montana Department of Environmental Quality (MT DEQ). Sampling consisted of D-frame net sweeps through emergent vegetation (when present), the water column, over the water surface, and included disturbing and scraping substrates at each sampled sites. Samples were preserved in ethanol at each wetland site and subsequently delivered to Rhithron Associates, Inc. for processing, taxonomic determinations, and data analysis.

At Rhithron's laboratory, Caton subsamplers and stereomicroscopes with 10X magnification were used to randomly select a minimum of 100 organisms, when possible, from each sample. In some cases, the entire sample contained fewer than 100 organisms; in these cases, all organisms from the sample were taken. Taxa were identified in general accordance with the taxonomic resolution standards set out in the MT DEQ Standard Operating Procedures for Sampling and Sample Analysis (Bukantis 1998). All samples were re-identified by a second taxonomist for quality assurance purposes. The identified samples have been archived at Rhithron's laboratory. Taxonomic data and organism counts were entered into an Excel 2000 spreadsheet, and metrics were calculated and scored using spreadsheet formulae.

Bioassessment metrics

An index based on the performance of 12 metrics was constructed, as described above. Table 1 lists those metrics, describes their calculation and the expected response of each to increased degradation or impairment of the wetland.

In addition to the summed scores of each metric and the associated impairment classification described above, each individual metric informs the bioassessment to some degree. The four richness metrics (Total taxa, POET, Chironomidae taxa, and Crustacea taxa + Mollusca taxa) can be interpreted to express habitat complexity as well as water quality. Complex, diverse habitats consist of variable substrates, emergent vegetation, variable water depths and other factors, and are potential features of long-established stable wetlands with minimal human disturbance. In the study conducted by Stribling et al. (1995), all four richness metrics were found to be significantly associated with water quality parameters including conductance, salinity, and total dissolved solids.

Four composition metrics (%Chironomidae, %Orthoclaadiinae of Chironomidae, %Crustacea + %Mollusca, and %Amphipoda) measure the relative contributions of certain taxonomic groups that may have significant responses to habitat and/or water quality impacts. For example, amphipods have been demonstrated to increase in abundance in alkaline conditions. Short-lived, relatively mobile taxa such as chironomids dominate ephemeral environments; many are hemoglobin-bearers capable of tolerating de-oxygenated conditions.

Two tolerance metrics (the Hilsenhoff Biotic Index and %Dominant taxon) were included in the bioassessment battery. The HBI indicates the overall invertebrate assemblage tolerance to nutrient enrichment, warm water, and/or low dissolved oxygen conditions. The percent abundance of the dominant taxon has been demonstrated to be strongly associated with pH, conductance, salinity, total organic carbon, and total dissolved solids.

Two trophic measures (%Collector-gatherers and %Filterers) may be helpful in expressing functional integrity of the invertebrate assemblage, which can be impacted by poor water quality or habitat degradation. High proportions of filtering organisms suggest nutrient and/or organic enrichment, while abundant collectors suggest more positive functional conditions and well-developed wetland morphology. These organisms graze periphyton growing on stable surfaces such as macrophytes.

RESULTS

In 2001, 29 sites were sampled statewide. Nineteen of these sites were revisited in 2002, and 13 new sites were sampled. In 2003, 17 sites that had been visited in both 2001 and 2002 were re-sampled, and 11 sites sampled for the first time in 2001 were re-visited. In addition, 2 new sites were sampled. In 2004, 25 sites were re-visited, and 6 new sites were sampled. Thus, the 2004 database contains data for 122 sampling events at 50 unique sites. Table 2 summarizes sites and sampling years.

Metric scoring criteria were re-developed each year as new data was added. For 2004, all 122 records were utilized. Ranges of individual metrics, as well as median metric values remained remarkably consistent in each of the 4 years; minimal changes resulted from the addition of new data in 2004. The summary metric values and scores for the 2004 samples are given in Tables 3a-3d.

Literature cited

Bollman, W. 1998. Montana Valleys and Foothill Prairies Ecoregion. Master's Thesis. (M.S.) University of Montana. Missoula, Montana.

Bukantis, R. 1998. Rapid bioassessment macroinvertebrate protocols: Sampling and sample analysis SOP's. Working draft. Montana Department of Environmental Quality. Planning Prevention and Assistance Division. Helena, Montana.

Stribling, J.B., J. Lathrop-Davis, M.T. Barbour, J.S. White, and E.W. Leppo. 1995. Evaluation of environmental indicators for the wetlands of Montana: the multimetric approach using benthic macroinvertebrates. Report to the Montana Department of Health and Environmental Science. Helena, Montana.

Table 1. Aquatic invertebrate metrics employed in the MTDT mitigation wetland monitoring study, 2001- 2004.

Metric	Metric Calculation	Expected Response to Degradation or Impairment
Total taxa	Count of unique taxa identified to lowest recommended taxonomic level	Decrease
POET	Count unique Plecoptera, Trichoptera, Ephemeroptera, and Odonata taxa identified to lowest recommended taxonomic level	Decrease
Chironomidae taxa	Count unique midge taxa identified to lowest recommended taxonomic level	Decrease
Crustacea taxa + Mollusca taxa	Count unique Crustacea taxa and Mollusca taxa identified to lowest recommended taxonomic level	Decrease
% Chironomidae	Percent abundance of midges in the subsample	Increase
Orthocladiinae/Chironomidae	Number of individual midges in the sub-family Orthocladiinae / total number of midges in the subsample.	Decrease
%Amphipoda	Percent abundance of amphipods in the subsample	Increase
%Crustacea + %Mollusca	Percent abundance of crustaceans in the subsample plus percent abundance of molluscs in the subsample	Increase
HBI	Relative abundance of each taxon multiplied times that taxon's modified Hilsenhoff Biotic Index value. These numbers are summed over all taxa in the subsample.	Increase
%Dominant taxon	Percent abundance of the most abundant taxon in the subsample	Increase
%Collector-Gatherers	Percent abundance of organisms in the collector-gatherer functional group	Decrease
%Filterers	Percent abundance of organisms in the filterer functional group	Increase

Table 2. Montana Department of Transportation Mitigated Wetlands Monitoring Project sites. 2001 – 2004.

2001	2002	2003	2004
Beaverhead 1	Beaverhead 1	Beaverhead 1	Beaverhead 1
Beaverhead 2	Beaverhead 2		
Beaverhead 3	Beaverhead 3		Beaverhead 3
Beaverhead 4	Beaverhead 4	Beaverhead 4	
Beaverhead 5	Beaverhead 5	Beaverhead 5	Beaverhead 5
Beaverhead 6	Beaverhead 6	Beaverhead 6	Beaverhead 6
Big Sandy 1			
Big Sandy 2			
Big Sandy 3			
Big Sandy 4			
Johnson-Valier			
VIDA			
Cow Coulee	Cow Coulee	Cow Coulee	
Fourchette - Puffin	Fourchette - Puffin	Fourchette - Puffin	Fourchette - Puffin
Fourchette - Flashlight	Fourchette - Flashlight	Fourchette - Flashlight	Fourchette - Flashlight
Fourchette - Penguin	Fourchette - Penguin	Fourchette - Penguin	Fourchette - Penguin
Fourchette - Albatross	Fourchette - Albatross	Fourchette - Albatross	Fourchette - Albatross
Big Spring	Big Spring	Big Spring	Big Spring
Vince Ames			
Ryegate			
Lavinia			
Stillwater	Stillwater	Stillwater	Stillwater
Roundup	Roundup	Roundup	Roundup
Wigeon	Wigeon	Wigeon	Wigeon
Ridgeway	Ridgeway	Ridgeway	Ridgeway
Musgrave - Rest. 1			
Musgrave - Rest. 2			
Musgrave - Enh. 1			
Musgrave - Enh. 2			
	Hoskins Landing	Hoskins Landing	Hoskins Landing
	Peterson - 1	Peterson - 1	Peterson - 1
	Peterson - 2		Peterson - 2
	Peterson - 4	Peterson - 4	Peterson - 4
	Peterson - 5	Peterson - 5	Peterson - 5
	Jack Johnson - main	Jack Johnson - main	
	Jack Johnson - SW	Jack Johnson - SW	
	Creston	Creston	Creston
	Lawrence Park		
	Perry Ranch		
	SF Smith River	SF Smith River	SF Smith River
	Camp Creek	Camp Creek	Camp Creek
	Kleinschmidt	Kleinschmidt - pond	Kleinschmidt - pond
		Kleinschmidt - stream	Kleinschmidt - stream
		Ringling - Galt	
			Circle
			Cloud Ranch Pond
			Cloud Ranch Stream
			Colloid
			Jack Creek
			Norem

Table 3a.

	BEAVER HEAD #1	BEAVER HEAD #3	BEAVER HEAD #5	BEAVER HEAD #6	BIG SPRING CREEK	CIRCLE	CLOUD RANCH POND	CLOUD RANCH STREAM	COLLOID	CRESTON
Total taxa	27	12	21	18	25	16	16	20	8	18
POET	3	0	2	3	4	2	2	4	2	3
Chironomidae taxa	7	5	5	5	8	5	6	11	1	2
Crustacea + Mollusca	7	3	4	6	7	1	6	1	1	7
% Chironomidae	0.33636	0.18888	0.39285	0.57547	0.44329	0.55855	0.41666	0.84	0.09090	0.06087
Orthoclaadiinae/Chir	0.05405	0.35294	0.06818	0.36065	0.27907	0.69354	0.4	0.16666	0	0
%Amphipoda	0.03636	0	0.01785	0.05660	0.05154	0	0.00925	0	0	0
%Crustacea + %Mollusca	0.31818	0.73333	0.05357	0.12264	0.18556	0.03603	0.36111	0.01	0.09090	0.73913
HBI	7.97169	7.88888	8.36363	8.15789	7.61855	7.19090	7.32291	4.84	6	6.92173
%Dominant taxon	0.2	0.57777	0.23214	0.25471	0.23711	0.38738	0.13888	0.38	0.27272	0.37391
%Collector-Gatherers	0.40909	0.75555	0.51785	0.62264	0.78350	0.05405	0.67592	0.74	0.18181	0.29565
%Filterers	0.12727	0	0	0	0.01030	0.15315	0.09259	0.17	0	0.06087
Total taxa	5	1	5	3	5	3	3	3	1	3
POET	3	1	1	3	5	1	1	5	1	3
Chironomidae taxa	5	3	3	3	5	3	3	5	1	1
Crustacea + Mollusca	5	1	3	5	5	1	5	1	1	5
% Chironomidae	3	3	3	1	1	1	1	1	5	5
Orthoclaadiinae/Chir	1	3	1	3	3	5	3	1	1	1
%Amphipoda	5	5	5	3	3	5	5	5	5	5
%Crustacea + %Mollusca	5	1	5	5	5	5	3	5	5	1
HBI	1	1	1	1	1	3	3	5	5	3
%Dominant taxon	5	1	5	5	5	3	5	3	5	3
%Collector-Gatherers	1	3	3	3	3	1	3	3	1	1
%Filterers	1	3	3	3	3	1	1	1	3	1
	40	26	38	38	44	32	36	38	34	32
	0.666667	0.433333	0.633333	0.633333	0.733333	0.533333	0.6	0.633333	0.566667	0.533333
	sub-optimal	poor	sub-optimal	sub-optimal	optimal	sub-optimal	sub-optimal	sub-optimal	sub-optimal	sub-optimal

Table 3b.

	FOURCHETTE CREEK ALBATROSS RESERVOIR	FOURCHETTE CREEK FLASHLIGHT RESERVOIR	FOURCHETTE CREEK PENGUIN RESERVOIR	FOURCHETTE CREEK PUFFIN RESERVOIR	JACK CREEK	MDT CAMP CREEK	MDT HOSKINS LANDING	MDT KLEINSCHMIDT CREEK	MDT KLEINSCHMIDT POND
Total taxa	18	23	19	22	23	35	25	19	19
POET	3	5	4	3	5	12	4	4	6
Chironomidae taxa	6	9	6	4	8	14	4	6	4
Crustacea + Mollusca	3	4	5	8	7	1	6	2	4
% Chironomidae	0.135135	0.265306	0.066116	0.247934	0.352113	0.37963	0.036697	0.438776	0.047619
Orthoclaadiinae/Chir	0.2	0.346154	0.625	0.3	0.52	0.585366	0.5	0.627907	0.8
%Amphipoda	0.126126	0.336735	0.578512	0.041322	0.028169	0	0.018349	0.010204	0.009524
%Crustacea + %Mollusca	0.684685	0.387755	0.77686	0.371901	0.380282	0.111111	0.541284	0.061224	0.190476
HBI	7.972973	7.216495	7.7	6.950413	7.647059	4.570093	6.59633	6.561224	6.67619
%Dominant taxon	0.495495	0.336735	0.561983	0.140496	0.15493	0.111111	0.366972	0.316327	0.552381
%Collector-Gatherers	0.873874	0.816327	0.702479	0.38943	0.394366	0.416667	0.091743	0.683673	0.114286
%Filterers	0	0.010204	0.132231	0.008264	0.042254	0.12037	0.018349	0.153061	0.047619
Total taxa									
POET	3	5	3	5	5	5	5	3	3
Chironomidae taxa	3	5	5	3	5	5	5	5	5
Crustacea + Mollusca	3	5	3	3	5	5	3	3	3
% Chironomidae	1	3	3	5	5	1	5	1	3
Orthoclaadiinae/Chir	5	3	5	3	3	3	5	1	5
%Amphipoda	3	3	5	3	5	5	5	5	5
%Crustacea + %Mollusca	3	1	1	3	5	5	5	5	5
HBI	1	3	1	3	3	5	3	5	5
%Dominant taxon	1	3	1	3	1	5	5	5	5
%Collector-Gatherers	1	5	1	5	5	5	3	5	1
%Filterers	5	5	3	1	1	1	1	3	1
	3	3	1	3	3	1	3	1	3
	32	44	32	40	46	46	48	42	44
	0.533333	0.733333	0.533333	0.666667	0.766667	0.766667	0.8	0.7	0.733333
	sub-optimal	optimal	sub-optimal	optimal	optimal	optimal	optimal	optimal	optimal

Table 3d.

	ROUNDUP	SOUTH FORK SMITH RIVER	STILLWATER	WIGEON
Total taxa	9	20	23	16
POET	0	5	4	3
Chironomidae taxa	4	7	9	5
Crustacea + Mollusca	3	3	4	3
% Chironomidae	0.55	0.482143	0.466667	0.314815
Orthoclaadiinae/Chir	0.072727	0.055556	0.244898	0.647059
%Amphipoda	0	0.071429	0.12381	0.481481
%Crustacea + %Mollusca	0.42	0.116071	0.180952	0.574074
HBI	8.89	6.589286	6.47619	7.534653
%Dominant taxon	0.28	0.294643	0.133333	0.481481
%Collector-Gatherers	0.56	0.839286	0.628571	0.657407
%Filterers	0.14	0	0	0.083333
Total taxa				
POET	1	3	5	3
Chironomidae taxa	1	5	5	3
Crustacea + Mollusca	3	5	5	3
% Chironomidae	1	1	3	1
Orthoclaadiinae/Chir	1	1	1	3
%Amphipoda	1	1	3	5
%Crustacea + %Mollusca	5	3	3	1
HBI	3	5	5	3
%Dominant taxon	1	5	5	3
%Collector-Gatherers	5	5	5	3
%Filterers	3	5	3	3
	1	3	3	1
	26	42	46	32
	0.433333	0.7	0.766667	0.533333
	poor	optimal	optimal	Sub-optimal

Aquatic Invertebrate Taxonomic Data

Site Name BIG SPRING CREEK

Date Collected 8 / 5 / 2004

Order	Family	Taxon	Count	Percent	Unique	BI	FFG
		Ostracoda	3	3.09%	Yes	8	CG
Amphipoda		Copepoda	4	4.12%	Yes	8	CG
	Talitridae	<i>Hyaella</i>	5	5.15%	Yes	8	CG
Basommatophora		Physidae	1	1.03%	Yes	8	SC
	Planorbidae	<i>Helisoma</i>	2	2.06%	Yes	6	SC
Coleoptera		Halipidae	3	3.09%	Yes	5	PH
Diplostraca		Cladocera	1	1.03%	Yes	8	CF
Diptera	Ceratopogonidae	Ceratopogoninae	3	3.09%	Yes	6	PR
	Chironomidae	<i>Ablabesmyia</i>	1	1.03%	Yes	8	CG
		<i>Acricoctopus</i>	9	9.28%	Yes	10	CG
		<i>Chironomus</i>	3	3.09%	Yes	10	CG
		<i>Parachironomus</i>	2	2.06%	Yes	10	PR
		<i>Parakiefferiella</i>	2	2.06%	Yes	6	CG
		<i>Paratanytarsus</i>	23	23.71%	Yes	6	CG
		<i>Psectrocladius</i>	1	1.03%	Yes	8	CG
		<i>Pseudochironomus</i>	2	2.06%	Yes	5	CG
Ephemeroptera	Eaetidae	<i>Callibaetis</i>	10	10.31%	Yes	9	CG
Haplotaxida	Naididae	<i>Nais</i>	11	11.34%	Yes	8	CG
Heteroptera	Corixidae	Corixidae	1	1.03%	Yes	10	PH
		<i>Hesperocorixa</i>	2	2.06%	Yes	10	PH
	Notonectidae	<i>Notonecta</i>	1	1.03%	Yes	5	PR
Isopoda	Asellidae	<i>Caecidotea</i>	2	2.06%	Yes	8	CG
Odonata	Coenagrionidae	Coenagrionidae	3	3.09%	No	7	PR
		<i>Enallagma</i>	1	1.03%	Yes	7	PR
	Libellulidae	<i>Leucorrhinia</i>	1	1.03%	Yes	9	PR
Grand Total			97				

Aquatic Invertebrate Data Summary

Project ID: MDT04LW
STORET Station ID:
Station Name: BIG SPRING CREEK


Activity ID:
Sample Date: 8/5/2004

Sample type	
SUBSAMPLE TOTAL ORGANISMS	97
Portion of sample used	13.33%
Estimated number in total sample	728
Conversion factor	10.088
Estimated number in 1 square meter	978
Sampling effort	
Habitat type	
EPT abundance	10
Taxa richness	24
Number EPT taxa	1
Percent EPT	10.31%

DOMINANCE		
TAXON	ABUNDANCE	PERCENT
Paratanvartus	23	23.71%
Nais	11	11.34%
Callibaetis	10	10.31%
Acricotopus	9	9.28%
Hyalella	5	5.15%
SUBTOTAL 5 DOMINANTS		
	58	59.79%
Copepoda	4	4.12%
Ostracoda	3	3.09%
Coenagrionidae	3	3.09%
Haliphus	3	3.09%
Ceratopogoninae	3	3.09%
TOTAL DOMINANTS	74	76.29%

TAXONOMIC COMPOSITION				TAXONOMIC RATIOS			
GROUP	PERCENT	ABUNDANCE	#TAXA	METRIC	VALUE		
Non-insect taxa	29.90%	29	8	EPT/Chironomidae	0.23		
Odonata	5.15%	5	3	Baetidae/Ephemeroptera	1.00		
Ephemeroptera	10.31%	10	1	Hydropsychidae/Trichopt	#DIV/0!		
Plecoptera	0.00%	0	0				
Heteroptera	4.12%	4	3				
Megaloptera	0.00%	0	0				
Trichoptera	0.00%	0	0				
Lepidoptera	0.00%	0	0				
Coleoptera	3.09%	3	1				
Diptera	3.09%	3	1				
Chironomidae	44.33%	43	8				


TOLERANCE/CONDITION INDICES			
Community Tolerance Quotient (CTQa)			102.00
Hilsenhoff Biotic Index			7.62
DIVERSITY			
Shannon H (log)			4.30
Shannon H (log2)			2.98
Margalef D			5.24
Simpson D			0.09
P evenness			0.12
VOLITINISM			
TYPE	ABUNDANCE	# TAXA	PERCENT
Multivoltine	61	12	62.89%
Univoltine	33	12	34.02%
Semivoltine	3	1	3.09%




TAXA CHARACTERS			
	#TAXA		PERCENT
Tolerant	9		26.80%
Sensitive	0		0.00%
Clinger	0		0.00%

FUNCTIONAL COMPOSITION				FUNCTIONAL RATIOS			
GROUP	PERCENT	ABUNDANCE	#TAXA	METRIC	VALUE		
Predator	11.34%	11	6	Scraper/Filterer	3.00		
Parasite	0.00%	0	0	Scraper/Scraper + Filtere	0.75		
Gatherer	78.35%	76	13				
Filterer	1.03%	1	1				
Herbivore	0.00%	0	0				
Piercer	6.19%	6	3				
Scraper	3.09%	3	2				
Shredder	0.00%	0	0				
Omnivore	0.00%	0	0				
Unknown	0.00%	0	0				

BIOASSESSMENT INDICES			
B-IBI (Karr et al.)			
METRIC	VALUE		SCORE
Taxa richness	24		3
E richness	1		1
P richness	0		1
T richness	0		1
Long-lived	1		1
Sensitive richness	0		1
%tolerant	26.80%		3
%predators	11.34%		3
Clinger richness	0		1
%dominance (3)	45.36%		5
TOTAL SCORE		20	40%



MONTANA DEQ INDICES (Bukantis 1998)				
METRIC	VALUE	Plains Ecoregions	Valleys and Foothills	Mountain Ecoregions
Taxa richness	24	2	2	2
EPT richness	1	0	0	0
Biotic Index	7.62	0	0	0
%Dominant taxon	23.71%	3	3	3
%Collectors	79.38%	2	1	1
%EPT	10.31%	1	0	0
Shannon Diversity	2.98	2		
%Scrapers +Shredder	3.09%	1	0	0
Predator taxa	6	3		
%Multivoltine	62.89%	1		
%H of T	#DIV/0!		#DIV/0!	
TOTAL SCORES		15	#DIV/0!	6
PERCENT OF MAXIMUM		50.00	#DIV/0!	28.57
IMPAIRMENT CLASS		MODERATE	#DIV/0!	MODERATE



COMMUNITY TOLERANCES	
Sediment tolerant taxa	0
Percent sediment tolerant	0.00%
Sediment sensitive taxa	0
Percent sediment sensitive	0.00%
Metals tolerance index (McGuire)	2.53
Cold stenotherm taxa	0
Percent cold stenotherms	0.00%

Montana Valleys and Foothills revised index (Bollman 1998)			
Percent max.	22.22%	Impairment class	MODERATE
Montana Plains ecoregions metrics (Bramblett and Johnson 2002)			
Riffle		Pool	
EPT richness	1	E richness	1
Percent EPT	10.31%	T richness	0
Percent Oligochaetes and Leeches	11.34%	Percent EPT	10.31%
Percent 2 dominants	35.05%	Percent non-insect	29.90%
Filterer richness	1	Filterer richness	1
Percent intolerant	0.00%	Univoltine richness	12
Univoltine richness	12	Percent supertolerant	58.76%
Percent clingers	0.00%		
Swimmer richness	5		

HABITUS MEASURES	
Hemoglobin bearer richness	4
Percent hemoglobin bearers	8.25%
Air-breather richness	0
Percent air-breathers	0.00%
Burrower richness	3
Percent burrowers	8.25%
Swimmer richness	5
Percent swimmers	17.53%